



Revista Mexicana de Ingeniería Química

ISSN: 1665-2738

amidiq@xanum.uam.mx

Universidad Autónoma Metropolitana

Unidad Iztapalapa

México

González-Blanco, G.; Valencia-Nava, A.; Beristain-Cardoso, R.; Hernández-Jaimes, C.;
Orozco-Villafuerte, J.; Buendía-González, L.

ANAMMOX ACTIVITY OF SLUDGE COMING FROM WETLAND MONOCOTS (*Typha*
sp.): KINETIC STUDY ACTIVIDAD ANAMMOX

Revista Mexicana de Ingeniería Química, vol. 16, núm. 2, 2017, pp. 555-561

Universidad Autónoma Metropolitana Unidad Iztapalapa

Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=62052087020>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

**ANAMMOX ACTIVITY OF SLUDGE COMING FROM WETLAND MONOCOTS
(*Typha* sp.): KINETIC STUDY****ACTIVIDAD ANAMMOX EN LOS LODOS PROCEDENTES DE
MONOCOTILEDÓNEAS DE HUMEDALES (*Typha* sp.): ESTUDIO CINÉTICO**

G. González-Blanco¹, A. Valencia-Nava¹, R. Beristain-Cardoso², C. Hernández-Jaimes¹,
J. Orozco-Villafuerte^{3*}, L. Buendía-González¹

¹Facultad de Ciencias, ³Facultad de Química, Universidad Autónoma del Estado de México, El Cerrillo, Piedras Blancas
Carretera Toluca-Ixtlahuaca km 15.5, Toluca, Edo de Méx., C.P. 50200.

²Departamento de Recursos de la Tierra, Universidad Autónoma Metropolitana-Unidad Lerma. Av. Hidalgo Pte. 46, Col.
Estación, Lerma de Villada, Edo de Méx., C.P. 52006.

Received September 6, 2016; Accepted January 12, 2017

Abstract

The anaerobic ammonium oxidation (anammox) activity was evaluated using sludge obtained from the rhizome of *Typha* sp., found in a natural wetland localized in the José Antonio Álzate dam, Toluca Valley, Mexico. Five NO_2^- -N concentrations (15, 30, 50, 70 and 100 mg/L) were evaluated and the ammonium remained constant at 100 ± 10 mg N/L. All cultures presented a lag phase in the ammonium consumption. Nevertheless, after the phase lag the ammonium oxidation was linked to nitrite reduction, producing N_2 . The nitrite consumption efficiencies for the concentrations evaluated from 15 to 50 mg/L of NO_2^- -N were above 95%, while the ammonium consumption efficiency increased from 14 to 48%. In experiments with 70 and 100 mg/L of NO_2^- -N, nitrite consumption efficiencies were 74 and 58% respectively. The higher nitrite concentrations tested inhibited the anammox process since the specific rates diminished at 0.51 and 0.44 mg NO_2^- -N/g VSS d respectively. This study provides new and useful information about the anammox activity in sludges coming from wetlands, and these can be used as potential inoculum to treat wastewaters into anammox reactors, or in constructed wetlands.

Keywords: anammox, wetlands, specific activity, nitrite, *Typha*.

Resumen


Se evaluó la actividad de oxidación anaerobia de amonio (anammox) usando un lodo obtenido del rizoma de *Typha* sp., de un humedal natural localizado en la presa José Antonio Álzate, en el valle de Toluca, México. Se evaluaron cinco concentraciones de N-NO_2^- (15, 30, 50, 70 y 100 mg / L) manteniendo constante la concentración de N-NH_4^+ en 100 ± 10 mg/L. Todos los cultivos presentaron una fase lag en el consumo de amonio, concluida esta fase, la oxidación del amonio estuvo acoplada a la reducción del nitrito, con producción de N_2 . Las eficiencias de consumo de nitrito para las concentraciones de 15 a 50 mg/L de N-NO_2^- fueron mayores a 95%, mientras que las eficiencias de amonio aumentaron de 14 a 48%. En los experimentos con 70 y 100 mg/L de N-NO_2^- , el consumo de nitrito fue de 74 y 58%, respectivamente observando un fenómeno de inhibición, reflejado en la disminución de las velocidades específicas (0.51 a 0.44 mg N-NO_2^- /g SSV d). El estudio proporciona información nueva y útil sobre la actividad anammox en lodos procedentes de humedales, los cuales pueden ser utilizados como inóculo potencial para el tratamiento de aguas residuales en reactores anammox ó en humedales construidos.

Palabras clave: anammox, humedales, actividad específica, nitrito, *Typha*.

* Corresponding author. E-mail: jov202001@yahoo.com.mx
Tel. 01-722-217-51-09; 217-38-90

1 Introduction

The Anaerobic Ammonia Oxidation (anammox) process was discovered in a denitrifying pilot plant (Mulder *et al.*, 1995). The anammox reaction involves the oxidation of ammonium coupled to the reduction of nitrite to produce the main product, molecular nitrogen, and a minor product, nitrate (Eq. 1) (Strous *et al.*, 1998) under anoxic conditions. The biological reaction is catalyzed by chemolithoautotrophic bacteria which have been identified as a distinct phylogenetic order, the Brocadiales, which is part of the phylum Planctomycetes (Kartal *et al.*, 2008; Jetten *et al.*, 2009). Since its discovery, it has gained attention due to its importance in the global nitrogen cycle that has been estimated accounts for over 50% of nitrogen loss in marine ecosystems (den Camp *et al.*, 2006). Anammox has mainly been investigated in wastewater treatment plants, enrichment cultures, in sediments and water columns (Dalsgaard *et al.*, 2003; Zhu *et al.*, 2013). However, there is few information of anammox in natural and constructed freshwater wetlands (Zhu *et al.*, 2010).



The José Antonio Alzate Dam (JAAD) in the Toluca Valley, Mexico, receives wastewaters of origin domestic and industrial, through the Lerma River (Barceló-Quintal *et al.*, 2012a, 2012b). The uncontrolled wastewaters discharges from municipal and industrial activities, as well as from agricultural runoffs, untreated or partially treated; induce contamination of water sources, risking more and more its complete use becoming a potential source of diseases. The main pollutants contained in the dam are organic matter, metals, phosphorous and reactive nitrogen (3 at 60 mg/L) (López-Galván *et al.*, 2010; Barceló-Quintal *et al.*, 2012a). Nitrogen removal is necessary because of the environmental impact, such as eutrophication, acidification and toxicity to aquatic life (Télliez-Pérez *et al.*, 2013). The dam have wetlands with monocots (*Typha*, *Juncus* and *Eichhornia*), the wetlands are buffer zones between land and waters that maintain good water quality for reducing nutrients and pollutants (Fisher and Acreman, 2004; Ramírez-Carrillo *et al.*, 2009; Trepel 2010; Wang and Gu, 2013). It has been shown in several studies the capacity of wetlands to remove organic matter, nitrogen, phosphorus, suspended solids, pathogens and heavy metals (Fisher and Acreman, 2004; Humbert *et al.*, 2012; Waki *et al.*, 2015). This capability is attributed to biological diversity they possess, for instance, it has been reported that the mechanism of nitrogen removal was conducted by: 1) ammonium assimilation by plants and 2) microbial activity through nitrification followed by denitrification (Verhoeven and Meuleman, 1999; Vymazal, 2007; Knox *et al.*, 2008; Lee *et al.*, 2009). However, the participation of anammox microorganisms, can also be assumed to contribute to nitrogen removal, but there is relatively little literature about the anammox activity in natural and constructed wetlands (Zhu *et al.*, 2010).

Most of the studies that have been conducted were focused on evaluating the diversity of anammox bacteria, abundance and population dynamics (Zhu *et al.*, 2011; Humbert *et al.*, 2012), distribution patterns (Waki *et al.*, 2015) but knowledge about the specific activity of anammox bacteria is scarce and particularly in soils wetland. Therefore, the goal of this study was focused on the respiratory activity of anammox process using anoxic sludge as inoculum potential in the treatment of dam's water. So in this paper the anammox activity was evaluated using sludge taken from the rhizome of *Typha* sp. of this natural wetland in the dam Jose Antonio Alzate from the Lerma River.

Most of the studies that have been conducted were focused on evaluating the diversity of anammox bacteria, abundance and population dynamics (Zhu *et al.*, 2011; Humbert *et al.*, 2012), distribution patterns (Waki *et al.*, 2015) but knowledge about the specific activity of anammox bacteria is scarce and particularly in soils wetland. Therefore, the goal of this study was focused on the respiratory activity of anammox process using anoxic sludge as inoculum potential in the treatment of dam's water. So in this paper the anammox activity was evaluated using sludge taken from the rhizome of *Typha* sp. of this natural wetland in the dam Jose Antonio Alzate from the Lerma River.

2 Materials and methods

2.1 Collection and preservation of microbial consortium

Ten plants of *Typha* sp. were collected of five sample points of the natural wetland present in the dam. It was collected with a portion of soil at the roots (Leander, 1972). These were carried at room temperature in plastic bags. The sludge obtained was spiked in an Erlenmeyer flask with 1000 mL of sterile isotonic solution (sodium chloride 0.90%). The mixing was homogenized at 20 min., and it allowed settling the sludge in an Imhoff cone. This operation was performed three times in order to determine the amount of total suspended solids (SSV). The biomass concentration quantified was 4.7 ± 0.2 g VSS/L (APHA, 2005).

2.2 Batch cultures

Batch cultures were conducted in 125 mL serological bottles containing 60 mL of basal medium formulated with the following compounds (mg/L): NaHCO_3

(2500); NaH₂PO₄·H₂O (57.5); CaCl₂·2H₂O (100); MgSO₄·7H₂O (200); and 1.5 mL/L of trace element solution. Trace element solution contained (in mg/L) FeSO₄ (5000) and ethylenediamine-tetraacetic acid (EDTA) (5000); ZnSO₄·7H₂O (430); CoCl₂·6H₂O (240); MnCl₂ (629); CuSO₄·5H₂O (250); Na₂MoO₄·2H₂O (220); NiCl₂·6H₂O (190); MgCl₂ (500). The initial sludge concentration was 2 ± 0.01 g of VSS/L the liquid and headspace were flushed during 3 min, with He/CO₂ mixture (80/20, v/v) to exclude oxygen from the assays. All bottles were sealed with butyl rubber stoppers and aluminum crimp seals. Separate bottles were set up for gas phase measurements of N₂. Abiotic controls lacking inoculum were run in parallel to monitor the possible abiotic elimination of the substrates. Biotic controls lacking nitrite or ammonium were also included to correct for compound losses not associated with anammox. All assays either abiotic or biotic were carried out by duplicate in an orbital shaker 150 rpm and 35°C. The initial pH for all batch cultures was of 7.5 ± 0.2. Five concentrations of N-NO₂⁻ (15, 30, 50, 70 and 100 mg/L) were evaluated and the concentration of electron donor remained constant at 100 ± 10 mg N-NH₄⁺/L. Microbial activity was evaluated in terms of consumption efficiency (Eq. 2), production yield (Eq. 3) and a specific consumption rate (Eq. 4). The kinetic parameters such as the specific rates were calculated using the Gompertz model and non-linear regression program (OriginPro 8.0). Each bottle was an independent experimental unit, which was sacrificed after sampling. Samples were filtered (0.45 μm) and analyzed for ammonium, nitrite, and nitrate determination.

$$\%E_N = \frac{mg\ N\ initial/L - mg\ N\ final/L}{mg\ N\ initial/L} \times 100 \quad (2)$$

$$Y_{P/S} = \frac{mg\ of\ product/L}{mg\ substrate\ consumed/L} \quad (3)$$

$$q_{N-NO_2^-} = \frac{mg\ NO_2^- - N\ consumed}{g\ VSS\ h} \quad (4)$$

2.3 Analytical methods

Ammonium was analyzed by a selective electrode (Phoenix electrode company, USA). Nitrite and nitrate were measured by ultraviolet spectrophotometry (Thermo Scientific, GENESYS 10S) with xenon lamp provides high intensity at 300 and 350 respectively and quartz cells 1 cm optical path (Modified of APHA 2005). Standard curves of nitrate and nitrite were prepared with the mineral medium used in batch

cultures to reduce interferences. Before sampling, all liquid samples were filtered through a 0.45-μm nylon membrane. Biogas production was measured using columns with NaCl solution (300 g/L, pH = 2) and the composition was determined only at the end of the kinetic by gas chromatography with thermal conductivity detector (Gow Mac model 550). Temperatures for the column, injector, and detector were 50, 100, and 110 °C, respectively. Helium was used as carrier gas at a constant flow rate of 16 mL/min. The stainless steel column (Porapak Q mesh 100-80) was of 1.20-m long and 1/8" diameter.

3 Results and discussion

3.1 Control assays

Batch assays were performed for a period of approximately 55 h. Abiotic cultures spiked with NO₂⁻-N and NH₄⁺-N did not show any chemical reaction, because at the end of the batch experiments the initial concentrations remained without change. Regarding to inoculated treatments, NO₂⁻-N consumption without electron donor (NH₄⁺-N) and vice versa were evaluated in order to observe contribution of a possible endogenous metabolism. At the end of the batch assays ammonium was almost completely recovered (~98 %), while for the nitrite assays it showed a consumption of 8%, this can be attributed to the endogenous metabolism or the presence of residual organics in the culture favoring the denitrification process (Peng and Zhu, 2006; Torà *et al.*, 2011). Finally, these experimental data showed that consumption of both substrates were mainly associated to the anammox activity of the sludge.

3.2 Anammox activity assays

Figure 1 shows the time-course of nitrogen compounds consumption. In batch cultures spiked with 15 mg/L NO₂⁻-N, it presented a lag phase of 2 h; while batch cultures with 70 and 100 mg/L NO₂⁻-N displayed lag phases around 5 h. Nonetheless, nitrite consumption did not present lag phase. Nitrite was consumed from the beginning, due to the endogenous metabolism as was described above. In overall, in all batch cultures the ammonium oxidation was linked to nitrite reduction, being the main product, N₂. For instance, the nitrite consumption efficiencies for the concentrations evaluated from 15 to 50 mg/L of NO₂⁻-N were above 95% (Table 1).




Fig. 1. Profiles of consume of an anammox enrichment culture to different concentrations of nitrite evaluated (A) NO_2^- -N; (B) NH_4^+ -N.

The ammonium consumption efficiency increased from 14 to 48%, this means, that initial nitrite concentration improved the ammonium consumption. The values obtained for the N_2 production yield indicated that total nitrogen consumed was recovered mainly as molecular nitrogen. The low efficiencies for ammonium consumption might suggest that initial nitrite concentration was not enough to convert all ammonium up to N_2 .

In batch assays evaluated with 70 and 100 mg/L of NO_2^- -N, nitrite consumption efficiencies were 74 and 58% respectively. The ammonium consumption efficiencies were 46 and 55%, respectively. At these

initial nitrite concentrations the substrate affected was the nitrite profile, due to ammonium consumption continued increasing. These low efficiencies might be associated to inhibitory troubles. For example, there are works indicating that anammox process can be inhibited by their substrates (nitrite and ammonium) and also for the nitrate produced. Nitrite is the most important inhibitor, because it has reported an IC_{50} for the nitrite specific activity ranging from 98 to 350 NO_2^- -N mg/L. Rowe *et al.* (1979) and Rake and Eagon (1980) suggested that nitrite is an uncoupler of the respiratory chain. In the case of ammonium, the IC_{50} reported ranges from 770-980 NH_4^+ -N mg/L and for nitrate of 630-980 NO_3^- -N mg/L (Strous *et al.*, 1999; Dapena-Mora *et al.*, 2007). The level of inhibition is strongly related to several factors such as cell concentration in the cultures, history of inoculum, among others.

The diminishing in the specific consumption rates for nitrite indicated inhibitory effects, as can be seen in the Figure 2. The last initial nitrite concentration evaluated was in the range of inhibition reported by Dapena-Mora *et al.*, 2007. At concentration of 100 mg/L of NO_2^- -N, nitrite affected stronger the respiratory process of anammox, since nitrite consumption diminished, and also the N_2 production diminished up to 0.81; this value suggested that 19% of total nitrogen consumed followed another metabolic fate. The nitrogen missing might be in form of anammox intermediates like hydroxylamine or hydrazine (Jetten *et al.*, 1999).

According to Sun *et al.* (2011), the activity of anammox process can be corroborated from the stoichiometry of the process. Strous *et al.*, (1998) established the stoichiometry of the anammox reaction, as can be seen in Eq. (1). The stoichiometric molar ratios in the anammox assays of this study were consistent with the stoichiometry of anammox reaction regardless of initial concentration of nitrite.

Table 1. Efficiencies consumption, performance N_2 and specific consumption rates nitrite at different concentrations of NO_2^- -N evaluated.

NO_2^- -N (mg /L)	$\%E_N \text{NO}_2^-$ -N	$\%E_N \text{NH}_4^+$ -N	Y_{N_2}	$q \text{NO}_2^-$ -N*	R^2
15.2 ± 0.6	99.6 ± 1.1	14.5 ± 0.8	0.98 ± 0.06	0.54 ± 0.03	0.98
30.3 ± 0.7	96.7 ± 0.9	28.2 ± 0.4	0.96 ± 0.05	0.56 ± 0.02	0.98
50.1 ± 0.5	97.3 ± 1.7	48.7 ± 0.3	0.98 ± 0.04	0.59 ± 0.04	0.91
70.2 ± 0.6	74.7 ± 1.2	46.3 ± 0.6	0.94 ± 0.06	0.51 ± 0.03	0.92
100.0 ± 0.6	58.9 ± 1.2	55.7 ± 1.2	0.81 ± 0.03	0.44 ± 0.03	0.94

qs * specific substrate consumption rate obtained by adjusting Gompertz

Table 2. Nitrogen balance, stoichiometric ratio of culture to different concentrations of nitrite evaluated

NO ₂ ⁻ -N Initial concentration (mg/L)	NH ₄ ⁺ -N Consumptions (mg/L)	NO ₂ ⁻ -N Consumptions (mg/L)	NO ₂ ⁻ removed/ NH ₄ ⁺ consumed Relations (mol/mol)
15.2 ± 0.6	10.5 ± 0.7	15.1 ± 0.2	1.39 ± 0.3
30.3 ± 0.7	21.1 ± 0.5	29.3 ± 0.8	1.38 ± 0.2
50.1 ± 0.5	36.5 ± 0.5	48.7 ± 1.1	1.33 ± 0.4
70.2 ± 0.6	34.9 ± 1.2	52.4 ± 0.9	1.51 ± 0.3
100.0 ± 0.6	41.7 ± 0.9	58.3 ± 0.5	1.40 ± 0.3




Fig. 2. NO₂⁻-N consumption rate evaluated in batch cultures. Batch cultures were carried out in duplicate. Error bars represent the standard deviation.

The reactant ratios of NO₂⁻ consumption to NH₄⁺ consumption were all very near the benchmark value of anammox stoichiometry at 1.32 (Table 2). Overall, the results indicated that anammox was the dominant process in the cultures.

Finally, the sludge taken from the roots of *Typha* sp. showed the metabolic capability for carrying out the anammox process. In this work, the characterization of the bacterial microflora was not possible. Nonetheless, phylogenetic studies in natural and artificial wetlands have identified anammox bacterial like *Candidatus* and *Anammoxoglobus* genus (Zhu et al., 2011; Waki et al., 2015).

Conclusions

The sludge taken from the roots of *Typha* sp. showed the potential to carry out the anammox biological process. At initial nitrite concentrations below to 70 mg/L were obtained high consumption efficiencies for nitrite, whereas for above, the consumption efficiencies for nitrite diminished. At higher nitrite concentrations was observed an inhibitory phenomenon, however, the reaction time was relatively short. Therefore, the use of this

kind of sludge might shorten the start-up process. The anaerobic ammonium oxidation is an emerging technology for nitrogen removal that provides a more environmentally sustainable and cost effective alternative compared to conventional biological treatments.

Nomenclature

<i>Anammox</i>	Anaerobic Ammonia Oxidation
% <i>E_N</i>	efficiency of consume of NO ₂ ⁻ -N or NH ₄ ⁺ -N, %
<i>JAAD</i>	Jóse Antonio Álzate dam
<i>q_{NO₂-N}</i>	specific substrate consumption rate mg of N consumed/g VSS h
<i>VSS</i>	volatile suspended solids, g/L
<i>Y_{P/S}</i>	anammox yield, mg of N ₂ produced/mg of N consumed

Acknowledgements

The first author acknowledges the support of the National Council for Science and Technology (CONACyT) through a scholarship to attend the post-doctorate within the Postgraduate Sciences of the Autonomous University of the State of Mexico, State of Mexico.

References

- American Public Health Association (APHA) (2005). *Standard Methods for the Examination of Water and Wastewater*, Washington DC, USA. 21 ed.
- Barceló-Quintal, I., López-Galván, E., Solís-Correa, H., Domínguez-Mariani, E., and Gómez-Salazar, S. (2012a). Water quality assessment of José Antonio Alzate dam, the Lerma River and its tributaries in the State of Mexico, Mexico.

- Journal of Environmental Protection* 3, 878-888.
- Barceló-Quintal, I.D., Solís-Correa, H.E., Avila-Pérez, P., López-Galván, E., Gómez-Salazar, S., and García-Albortante, J. (2012b). Determination of distributions of Cd, Cu, and Pb concentrations in sediments of a Mexican reservoir to infer their environmental risk. *Biological Trace Element Research* 148, 122-132.
- Dalsgaard, T., Canfield, D.E., Petersen, J., Thamdrup, B. and Acuña-González, J. (2003). N₂ production by the anammox reaction in the anoxic water column of Golfo Dulce, Costa Rica. *Nature* 422, 606-608.
- Dapena-Mora, A., Fernández, I., Campos, J.L., Mosquera-Corral, A., Méndez, R. and Jetten, M.S.M. (2007). Evaluation of activity and inhibition effects on anammox process by batch tests based on the nitrogen gas production. *Enzyme and Microbial Technology* 40, 859-865.
- den Camp, H., Kartal, B., Guven, D., van Niftrik, L., Haaijer, S.C.M., van der Star, W.R.L., van de Pas-Schoonen, K.T., Cabezas, A., Ying, Z., Schmid, M.C., Kuypers, M.M.M., van de Vossenberg, J., Harhangi, H.R., Picioreanu, C., van Loosdrecht, M.C.M., Kuenen, J.G., Strous, M. and Jetten, M.S.M. (2006). Global impact and application of the anaerobic ammonium-oxidizing (anammox) bacteria. *Biochemical Society Transactions* 34, 174-178.
- Fisher, J. and Acreman, M. C. (2004). Wetland nutrient removal: a review of the evidence. *Hydrology and Earth System Sciences Discussions* 8, 673-685.
- Humbert, S., Zopfi, J., Tarnawski, S. E. (2012). Abundance of anammox bacteria in different wetland soils. *Environmental Microbiology Reports* 4, 484-490.
- Jetten, M.S.M., Strous, M., van de Pas-Schoonen, K.T., Schalk, J., van Dongen, L., van de Graaf, A.A., Logemann, S., Muyzer, G., van Loosdrecht, M.C.M. and Kuenen, J.G. (1999). The anaerobic oxidation of ammonium. *FEMS Microbiology Review* 22, 421-437.
- Jetten, M.S.M., Niftrik, L., Strous, M., Kartal, B., Keltjens, J.T. and Op den Camp, H.J. (2009). Biochemistry and molecular biology of anammox bacteria. *Critical Reviews in Biochemistry and Molecular Biology* 44, 65-84.
- Kartal, B., Van Niftrik, L., Rattray J., Van De Vossenberg, J.L.C.M., Schmid, M.C., Sinninghe-Damste, J., Jetten, M.S.M. and Strous, M. (2008). Candidatus Brocadia fulgida: An autofluorescent anaerobic ammonium oxidizing bacterium. *FEMS Microbiology Ecology* 63, 46-55.
- Knox, A.K., Dahlgren, R.A., Tate, K.W. and Atwill, E.R. (2008). Efficacy of natural wetlands to retain nutrient, sediment and microbial pollutants. *Journal of Environmental Quality* 37, 1837-1846.
- Kuypers, M.M., Sliemers, A.O., Lavik, G., Schmid, M., Jørgensen, B.B., Kuenen, J.G., ...and Jetten, M. S. (2003). Anaerobic ammonium oxidation by anammox bacteria in the Black Sea. *Nature* 422, 608-611.
- Lee, C.G., Fletcher, T.D. and Sun, G. (2009). Nitrogen removal in constructed wetland systems. *Engineering in Life Sciences* 9, 11-22.
- Leander, J. (1972). *Methods for Research on the Ecology of Soil-Borne Plant pathogens*, Ed. Burgess Publishing, USA, p.p. 37-57.
- López-Galván, E., Barceló-Quintal, I., Solís-Correa, H.E., Bussy, A.L., Avila-Pérez, P., and Delgadillo, S.M. (2010). Calculation of the Ecological Risk Index in the José Antonio Alzate Dam, State of Mexico, Mexico. *Biological Trace Element Research* 135, 121-135.
- Mulder, A., Graaf, A.A., Robertson, L.A. and Kuenen, J.G. (1995). Anaerobic ammonium oxidation discovered in a denitrifying fluidized bed reactor. *FEMS Microbiology Ecology* 16, 177-184.
- Peng, Y. and Zhu, G. (2006). Biological nitrogen removal with nitrification and denitrification via nitrite pathway. *Applied Microbiology and Biotechnology* 73, 15-26.
- Rake, J. B., Eagon, R. G. (1980). Inhibition, but not uncoupling, of respiratory energy coupling of three bacterial species by nitrite. *Journal of Bacteriology* 144, 975-982.

- Ramírez-Carrillo, H. F., Luna-Pabello, V. M., Arredondo-Figueroa, J. L. (2009). Evaluation of an intermittent artificial vertical flow wetland, to obtain good quality water for aquaculture. *Revista Mexicana de Ingeniería Química* 8, 93-99.
- Rowe, J.J., Yarbrough, J.M., Rake, J.B., Eagon, R.G. (1979). Nitrite inhibition of aerobic bacteria. *Current Microbiology* 2, 51-54.
- Strous, M., Heijnen, J.J., Kuenen, J.G. and Jetten, M.S.M. (1998). The sequencing batch reactor as a powerful tool for the study of slowly growing anaerobic ammoniumoxidizing microorganisms. *Applied Microbiology and Biotechnology* 50, 589-596.
- Strous, M., Fuerst, J.A., Kramer E.H.M., Logemann, S., Muyzer, G., Van de Pas-Schoonen, K.T., Webb, R., Kuenen, J.G. and Jetten, M.S.M. (1999). Missing lithotroph identified as new planctomycete. *Nature* 400, 446-449.
- Sun, W., Banihani, Q., Sierra-Alvarez, R., and Field, J. A. (2011). Stoichiometric and molecular evidence for the enrichment of anaerobic ammonium oxidizing bacteria from wastewater treatment plant sludge samples. *Chemosphere* 84, 1262-1269.
- Télliz-Pérez, S.K., Silva, C.D., and Texier, A.C. (2013). Simultaneous ammonium and p-hydroxybenzaldehyde oxidation in a sequencing batch reactor. *Revista Mexicana de Ingeniería Química* 12, 97-104.
- Thamdrup, B. and Dalsgaard, T. (2002). Production of N₂ through anaerobic ammonium oxidation coupled to nitrate reduction in marine sediments. *Applied Environmental and Microbiology* 68, 1312-1318.
- Torà, J.A., Baeza, J.A., Carrera, J. and Oleszkiewicz, J.A. (2011). Denitrification of a high-strength nitrite wastewater in a sequencing batch reactor using different organic carbon sources. *Chemical Engineering Journal* 172, 994-998.
- Trepel, M. (2010). Assessing the cost-effectiveness of the water purification function of wetlands for environmental planning. *Ecological Complexity* 7, 320-326.
- Verhoeven, J.T. and Meuleman, A.F. (1999). Wetlands for wastewater treatment: opportunities and limitations. *Ecological Engineering* 12, 5-12.
- Vymazal, J. (2007). Removal of nutrients in various types of constructed wetlands. *Science of the Total Environment* 380, 48-65.
- Waki, M., Yasuda, T., Suzuki, K., Komada, M. and Abe, K. (2015). Distribution of anammox bacteria in a free-water-surface constructed wetland with wild rice (*Zizania latifolia*). *Ecological Engineering* 81, 165-172.
- Wang, Y.F., and Gu, J.D. (2013). Higher diversity of ammonia/ammonium-oxidizing prokaryotes in constructed freshwater wetland than natural coastal marine wetland. *Applied Microbiology and Biotechnology* 97, 7015-7033.
- Zhu, G., Jetten, M.S.M., Kuschik, P., Ettwig, K.F. and Yin, C. (2010). Potential roles of anaerobic ammonium and methane oxidation in the nitrogen cycle of wetland ecosystems. *Applied Microbiology and Biotechnology* 86, 1043-1055.
- Zhu G, Wang S, Feng X, Fan G, Jetten MSM and Yin, C. (2011) Anammox bacterial abundance, biodiversity and activity in a constructed wetland. *Environmental Science and Technology* 45, 9951-9958.
- Zhu, G., Wang, S., Wang, W., Wang, Y., Zhou, L., Jiang, B., and Hefting, M. M. (2013). Hotspots of anaerobic ammonium oxidation at land-freshwater interfaces. *Nature Geoscience* 6, 103-107.