



OPEN ACCESS

EDITED BY

Vincenzo Auriemma,
University of Salerno, Italy

REVIEWED BY

Agata Benfante,
University of Turin, Italy
Serena Quarta,
University of Salerno, Italy

*CORRESPONDENCE

Mojtaba Habibi Asgarabad
✉ Mojtaba.h.asgarabad@ntnu.no
Zahra Vahabi
✉ zvahabi@sina.tums.ac.ir

SPECIALTY SECTION

This article was submitted to
Health Psychology,
a section of the journal
Frontiers in Psychology

RECEIVED 26 December 2022

ACCEPTED 30 March 2023

PUBLISHED 05 June 2023

CITATION

Habibi Asgarabad M, Etesam F, Salehi Yegaei P,
Vahabi Z, Akbari Saneh N, Fathi F, Ghosi F and
Wiiium N (2023) Measurement invariance of the
General Health Questionnaire (GHQ-12) across
gender and age: Demographic and medical
correlates of mental health in patients with
COVID-19.

Front. Psychol. 14:1132154.

doi: 10.3389/fpsyg.2023.1132154

COPYRIGHT

© 2023 Habibi Asgarabad, Etesam, Salehi Yegaei, Vahabi, Akbari Saneh, Fathi, Ghosi and Wiiium. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Measurement invariance of the General Health Questionnaire (GHQ-12) across gender and age: Demographic and medical correlates of mental health in patients with COVID-19

Mojtaba Habibi Asgarabad^{1,2,3,4,5*}, Farnaz Etesam^{6,7},
Pardis Salehi Yegaei¹, Zahra Vahabi^{6,8*}, Niusha Akbari Saneh⁹,
Fateme Fathi¹⁰, Fateme Ghosi¹¹ and Nora Wiiium¹²

¹Health Promotion Research Center, Iran University of Medical Science, Tehran, Iran, ²Department of Psychology, Norwegian University of Science and Technology, Trondheim, Norway, ³Department of Health Psychology, School of Behavioral Sciences and Mental Health, Tehran Psychiatric Institute, Iran University of Medical Sciences, Tehran, Iran, ⁴Positive Youth Development Lab, Human Development and Family Sciences, Texas Tech University, Lubbock, TX, United States, ⁵Center of Excellence in Cognitive Neuropsychology, Institute for Cognitive and Brain Sciences, Shahid Beheshti University, Tehran, Iran, ⁶Department of Geriatric Medicine, Tehran University of Medical Sciences, Ziaei Hospital, Tehran, Iran, ⁷Psychosomatic Research Center, Department of Psychiatry, Tehran University of Medical Sciences, Imam Khomeini Hospital, Tehran, Iran, ⁸Cognitive Neurology and Neuropsychiatry Division, Department of Psychiatry, Tehran University of Medical Sciences, Roozbeh Hospital, Tehran, Iran, ⁹Department of Clinical Psychology, School of Behavioral Sciences and Mental Health, Tehran Psychiatric Institute, Iran University of Medical Sciences, Tehran, Iran, ¹⁰Faculty of Psychology and Educational Sciences, Semnan University, Semnan, Iran, ¹¹Department of Psychology and Educational Sciences, University of Al Zahra, Tehran, Iran, ¹²Department of Psychosocial Science, Faculty of Psychology, University of Bergen, Bergen, Norway

Introduction: The present cross sectional study aimed to evaluate the construct and criterion validity, reliability, and gender and age differences of the 12-item General Health Questionnaire (GHQ-12) among hospitalized patients with COVID-19 in 2020. The criterion validity was assessed *via* its link with perceived stress, sleep quality, daily life activities, and demographic and medical characteristics.

Methods: A total of 328 COVID-19 patients (55.8% men; $M_{age} = 50.49$, $SD = 14.96$) completed the GHQ-12, the Perceived Stress Scale (PSS), the Pittsburgh Sleep Quality Index (PSQI), the Activities of Daily Life (ADL)-Katz Scale, and the Lawton Instrumental Activities of Daily Living Scale (IADL).

Results: Among 13 factorial models, the three-factor model (successful coping, self-esteem, and stress) was shown to have the best fit. GHQ-12 was positively associated with PSQI, PSS, Hyperlipidemia, psychiatry disorders, hospitalization duration, the change in sleep time, and use of sleeping pills, and negatively correlated with educational level, and the number of family members. The GHQ-12 also had a negative correlation with ADL and IADL in over 60 years of age group. Females scored higher on total GHQ-12 scores, compared to males. Finally, the hospitalization duration was longer for patients over 60 (mean = 8.8 days, $SD = 5.9$) than patients under 60 (mean = 6.35 days, $SD = 5.87$).

Discussion: Overall, the findings provided evidence that mental distress in patients with COVID-19 is correlated with high perceived stress, low sleep quality, low ADL and IADL, and a range of demographic features and medical conditions. Designing

psychological interventions for these patients that target the aforementioned correlates of mental distress is warranted.

KEYWORDS

COVID-19, concurrent validity, general health, medical condition, reliability, sleep, activities of daily life, stress

Introduction

New measures of the COVID-19 pandemic (e.g., self-isolation and quarantine) have led to an increase in mental distress, such as anxiety, insomnia, and suicidal attempt (WHO, 2020; Santomauro et al., 2021). Review studies after the COVID-19 pandemic showed high prevalence rate of anxiety (27–41.3%) and depression (27 to 34.1%) in Eastern Europe (Zhang et al., 2022), Southeast Asia (Pappa et al., 2022) and South Asia (Hossain et al., 2021).

To capture changes in mental health in both general and clinical populations, the self-administered General Health Questionnaire (GHQ; Goldberg, 1988) was developed. The 12-item short-form of GHQ (GHQ-12; Goldberg et al., 1997) was derived from the original 60-item questionnaire for fast administration in busy clinical settings. This questionnaire screens those with common psychological problems, such as poor self-esteem, stress, and sleep loss (del Pilar Sánchez-López and Dresch, 2008).

Although GHQ-12 was originally designed as a unidimensional measure (Goldberg, 1972), several exploratory factor analyses indicated that a two-factor (Zhong et al., 2021) or three-factor model (del Pilar Sánchez-López and Dresch, 2008) is the most common model (full information of these factorial models and their reliabilities are presented in Table 1). However, considering the study of Liang et al. (2016) for instance, they failed to find the best fitting model among ten existing factorial models, highlighting that there is a need for further research on the factor structure of GHQ-12.

General mental health: Stress and sleep quality

Mental health is reciprocally linked to sleep quality, specifically in patients with COVID-19 who are commonly susceptible to sleep disturbances (Deng et al., 2021; Marvaldi et al., 2021). Perceived stress, referring to the extent to which a person perceives their daily life situations as stressful, was also found to be positively associated with GHQ-12 in a large cross-national sample of COVID-19 patients (Bonsaksen et al., 2022). Patients infected with COVID-19 experienced the burden of job loss (Crayne, 2020), death anxiety

(Korkut, 2022), and may react with heightened stress (Bonsaksen et al., 2022).

General mental health: Medical conditions and demographic characteristics

Coronavirus may severely impair the subsequent physical functioning in some patients, especially the elderly (Carfi et al., 2020; Halpin et al., 2020). The Activities of daily living (ADL) and the Instrumental Activities of Daily Living (IADL; Katz et al., 1970; Lawton, 2000) are basic skills necessary for independently taking care of oneself (Edemekong et al., 2017) and environmental adaptation (Roehrig et al., 2007). Although few in number, some studies have shown the link of mental health to ADL and IADL scores (de Castro Costa et al., 2008).

Due to the negative effect of COVID-19 on several aspects of people's lives (Santomauro et al., 2021), research should document the association of mental health status in COVID-19 patients with demographics and medical features. Previous studies have suggested the positive link of mental health status with medical features, such as change in sleep time before and after COVID-19 and use of sleeping pills (Becker et al., 2018), hospitalization duration (Liao et al., 2020), psychiatry disorders (Kaufman et al., 2020), hyperlipidemia (HLP; Chang et al., 2021), diabetes (Moradian et al., 2021), cardiovascular disease (CVD; de Paiva Teixeira et al., 2020), substance use history (Czeisler et al., 2020), as well as demographics characteristics, namely, educational level (Dalgard et al., 2007), unemployment (Achdut and Refaeli, 2020), and the number of family members (Hendriksen et al., 2021). COVID-19 pandemic was shown to result in greater mental distress in women (Giorgi et al., 2014; Bucciarelli et al., 2021). As an example, in a study on large data of 49,156 participants, Proto and Quintana-Domeque (2021) found that after the COVID-19 pandemic, women manifested higher elevation in GHQ-12 scores (higher psychological distress) than men. Aging is believed to be associated with decrease in mental distress (Hoeymans et al., 2004), while COVID-19 pandemic have led to greater mental distress in younger patients (Bruine de Bruin, 2021).

The study context

Iran is among the worst-hit countries by Coronavirus, with heavy death tolls (more than 19,000 deaths until August 2020; Shahriarirad et al., 2021). Challenging factors, namely, the shortage of hygiene and medical supplies and equipment (i.e., masks and disinfectants), economic constraints (Zandifar and Badrfam, 2020), and the incapacity of government to formulate and enforce effectual social distancing and lockdown measures have led to the mental distress in the Iranian general population (Moghanibashi-Mansourieh, 2020; Zandifar and

Abbreviations: GHQ, General Health Questionnaire; PSS, Perceived Stress Scale; PSQI, Pittsburgh Sleep Quality Index; ADL, activities of daily life; IADL, instrumental activities of daily living; HTN, high blood pressure; HLP, hyperlipidemia; CVA, cerebrovascular accident; HPA, hypothalamic pituitary adrenal; PD, pulmonary disease; IDD, immune deficiency disease; MI, myocardial infarction; CVD, cardiovascular disease.

TABLE 1 Studies Validating the Psychometric Properties of the GHQ-12 in Different Populations.

Author	Country	Participant characteristics	Factor structure and fit indices	Factors and corresponding items	Reliability
Unidimensional models					
Alaminos-Torres et al. (2021)	Spain	n = 342 Age range = 41–50 years	Range of factor loadings (EFA; unidimensional model) = 0.57–0.95	–	Total score = 0.85
Gnams and Staufienbiel (2018)	Germany	n ₁ = 76,473 n ₂ = 410,640	Fit index (CFA; unidimensional model) = CFI = 0.89, RMSEA = 0.11	–	Total score = 0.85
Hystad and Johnsen (2020)	Norway	n ₁ = 591 n ₂ = 196	Fit index (CFA; unidimensional model) = CFI = 0.91, RMSEA = 0.07	–	–
Romppel et al. (2013)	Germany	n = 2,041 (53% female) M _{age} (SD) = 48.8 (18.1)	Fit indexes (CFA; unidimensional model) = CFI = 0.93, RMSEA = 0.10	–	Total score = 0.89 Positively worded items = 0.79 Negatively worded items = 0.86
Two-factor models					
Hamad (2022)	Saudi Arabia	n = 473 (60.81% female)	Fit index (CFA; two-factor model) = CFI = 0.96, RMSEA = 0.05	F1 Personal and Social Dysfunction = 1,3,4,7,8,9,10,11,12 F2 Anxiety = 2,5,6	Total score = 0.85
Kalliath et al. (2004)	New Zealand	n ₁ = 691 (54% female) M _{age} = 38 n ₂ = 415 (54% female) M _{age} = 38	Fit index (CFA; two-factor model) = CFI = 0.98, RMSEA = 0.07	F1 Social Dysfunction = 4,7,8,12 F2 Anxiety/ Depression = 6,9,10,11	Total score at T ₁ = 0.91 Total score at T ₂ = 0.90
Montazeri et al. (2003)	Iran	n = 748 (76% female) M _{age} (SD) = 21.1 (2.1)	Range of factor loadings (EFA; two-factor model) = F ₁ = 0.56–0.81; F ₂ = 0.46–0.69	F1 Psychological distress = 1,3,4,7,8,10,11 F2 social dysfunction = 2,5,6,7,9,12	Total score = 0.87
Najarkolaei et al. (2014)	Iran	n = 428 (56% female) M _{age} (SD) = 22.83 (3.09)	Range of factor loadings (EFA; two-factor model) = F ₁ = 0.46–0.73; F ₂ = 0.39–0.78 Fit indexes (CFA; two-factor model) = GFI = 0.96, RMSEA = 0.04	F1 social dysfunction = 1,2,5,7,9,12 F2 psychological distress = 3,4,6,8,10,11	Total score = 0.85 Social dysfunction = 0.77 Psychological distress = 0.76
Politi et al. (1994)	Italy	n = 320 (0% female) M _{age} (SD) = 18	Range of factor loadings (EFA; two-factor model) = F ₁ = 0.31–0.80; F ₂ = 0.34–0.72	F1 general dysphoria = 2,5,6,9,10,11,12 F2 social dysfunction = 1,3,4,7,8	Total score = 0.81
Schnitz et al. (1999)	Germany	n = 572 (68.7% female) M _{age} (SD) = 42.7 (15.7)	A principal-components factor analysis = factors' eigenvalues of >1	F1 Anxiety/ Depression = 1,2,6,7,10,11 F2 Social Performance = 4, 5, 8, 9 and 12	Anxiety/ Depression = 0.86 Social Performance = 0.82 Total score = 0.91
Zhong et al. (2021)	China	–	Range of factor loadings (EFA; two-factor model) = F ₁ = 0.62–0.72; F ₂ = 0.48–0.79 Fit indexes (CFA; two-factor model) = 0.92, RMSEA = 0.08	F1 = 1,2,5,7,9,12 F2 = 3,4,6,8,10,11	Total score = 0.89
Three-factor models					
Daradkeh et al. (2001)	United Arab Emirates	n = 157	Range of factor loadings (EFA; three-factor model) = F ₁ = 0.57–0.79; F ₂ = 0.44–0.80; F ₃ = 0.55–0.87	F1 general dysphoria = 10, 5, 9, 11, 6 F2 lack of enjoyment = 7, 12, 1, 8, 2 F3 social dysfunction = 3, 4	Total score = 0.86

(Continued)

TABLE 1 (Continued)

Author	Country	Participant characteristics	Factor structure and fit indices	Factors and corresponding items	Reliability
del Pilar Sánchez-López and Dresch (2008)	Spain	$n = 1,001$ (60% female) M_{age} (SD) = 41.75 (10.95)	Range of factor loadings (EFA; three-factor model) = $F_1 = 0.50-0.71$; $F_2 = 0.41-0.63$; $F_3 = 0.63-0.65$	F1 Successful Coping = 1,3,4,7,8,12 F2 Self-esteem = 6,9,10,11 F3 Stress = 2,5,9	Total score = 0.76
Farrell (1998)	Australia	$n = 270$ (85% female) M_{age} (SD) = 0.36 (9)	Range of factor loadings (EFA; three-factor model) = $F_1 = 0.62-0.81$; $F_2 = 0.60-0.80$; $F_3 = 0.70-0.84$	F1 Anxiety = 10,12,2,5,11 F2 Depression = 1,9,8,7,6 F3 Social dysfunction = 3,4	Anxiety = 0.84 Depression = 0.81 Social dysfunction = 0.69
Gao et al. (2004)	Singapore	$n = 120$ (47.5% female) M_{age} (SD) = 43.1 (12.7)	Fit index (CFA; three-factor model) = CFI = 0.93, RMSEA = 0.10	F1 Anxiety and depression = 2,5,9,6, F2 Social dysfunction = 1,3,4,8,7,12 F3 Loss of confidence = 10,11	–
Graetz (1991)	Australia	$n = 8,998$ (49% female) Age range = 16–25 years	Range of factor loadings (EFA; three-factor model) at T_1 and $T_2 = F_1 = 0.44-0.78$; $F_2 = 0.44-0.59$; $F_3 = -0.70 - -0.72$ Range of factor loadings (EFA; three-factor model) at T_3 and $T_4 = F_1 = 0.38-0.77$; $F_2 = 0.44-0.64$; $F_3 = -0.72 - -0.74$	F1 Anxiety and depression = 2,5,6,9 F2 Social dysfunction = 1,3,4,7,8,12 F3 loss of confidence = 10,11	–
Lee and Kim (2020)	South Korea	$n = 504$ (66.8% female) M_{age} (SD) = 20.2 (1.63)	Fit indexes (CFA; three-factor model) = CFI = 0.93, RMSEA = 0.07	F1 Anxiety and depression = 2,5,6,9 F2 Social dysfunction = 1,3,4,7,8,12 F3 loss of confidence = 10,11	Total score = 0.81
Liang et al. (2016)	China	$n = 1,051$ (38.5% female) Age range = 29–35 years	Three-dimensional model (CFA) = CFI = 0.98, RMSEA = 0.03	F1 = 4,6,9,10,11,12 F2 = 3,5,7,8 F3 = 1,2	Total score = 0.84
Martin (1999)	Australia	$n = 169$ (61.1% female) M_{age} (SD) = 28 (11)	Range of factor loadings (EFA; three-factor model) = $F_1 = 0.46-0.64$; $F_2 = 0.65-0.70$; $F_3 = -0.63-0.84$	F1 Self-esteem = 1,3,4,8 F2 Stress = 2,5,7 F3 Successful Coping = 6,9,10,11,12	Self-esteem = 0.83 Stress = 0.71 Successful Coping = 0.67

n , sample size; F , factor; T , study wave; M , mean; CFI, comparative fit index; RMSEA, root mean square error of approximation; GFI, goodness of fit index; EFA, exploratory factor analysis; CFA, confirmatory factor analysis. Inconsistency in reporting the demographic characteristics including sample size, percentage of females, $Mean_{age}$ (SD) in column “participants” is due to not reporting the relevant data in the papers.

Badrfam, 2020; Shahriarirad et al., 2021). The Iranian population has been no exception to the global trend of increased mental issues. In a group of 5,328 individuals from the general population of Iran, the prevalence rates of anxiety, depression, and comorbid depression-anxiety were determined to be 30.1, 33.4, and 22.1%, respectively (Nakhostin-Ansari et al., 2020). Moreover, in another recent study by Maroufizadeh et al. (2022), the prevalence of mild-to-severe anxiety and depression in Iranian medical students was found to be 38.1 and 27.6%, showing a significant impact on sleep patterns.

In Iran, two studies have assessed the psychometric properties of GHQ-12. In a study on emerging adults, Montazeri et al. (2003) findings confirmed the two-factor model, comprising “depression” and “social dysfunction.” Their study showed the negative association of GHQ-12 with global quality of life, supporting its satisfactory convergent validity. Similarly, the results of Najarkolaei et al. (2014) study supported a two-factor model including “distress” and “social dysfunction” in freshmen university students. Nevertheless, participants of these two studies had limited age range (18–26 years of age) and were recruited from university students, which prevent their results from being generalized to the general or clinical Iranian population.

The present study was first-of-its-kind that aimed to examine the psychometric properties of the GHQ-12 in Iranian patients with COVID-19. In specific, we aimed to examine: (1) the factor structure by conducting Confirmatory Factor Analysis (CFA) based on 13 empirically-derived factorial models, (2) the internal consistency, (3) the criterion validity through the relationship of GHQ with perceived stress, sleep quality, ADL/IADL, and demographic and medical variables, and (4) the comparison of the average GHQ-12 scores among age and gender groups (if any). We hypothesized that higher GHQ-12 score—that reflects lower mental health—has a positive relationship with: (1) poor sleep quality, (2) higher perceived stress, and (3) lower level of ADL/IADL functions.

Materials and methods

Participants

Participants comprised a total of 328 patients with COVID-19 (55.8% men), aged 21 to 92 ($Mean_{age}$ (SD) = 50.49 (14.96); 73.6%

60 years old or younger). As for educational level, 19.8% of participants were illiterate, 22.9% had primary education, 16.5% had secondary education, 23.5% had diploma level, and 17.3% had higher education. Their job status included 15.3% employee, 6.7% skill-worker, 20.7% self-employed, 39.6% unemployed, and 13.1% retired. Most patients (88.6%) were living with their spouse and/or their children, while 10.5% were living alone. A majority of 86.6% had no history of smoking, while 10.1 and 2.4% reported smoking in the past and at the present time. In addition, 93.9 and 96.6% reported no history of alcohol and drug use, respectively. Among participants, 4.6% reported using sleeping pills—mostly Alprazolam, Chlordiazepoxide, and Asentra. Patients under and over 60 reported underlying diseases, including Diabetes (13 and 44.6%), High Blood Pressure (HTN; 15.2 and 28.9%), HLP (11.7 and 31.3%), psychiatry disorders (4.8 and 3.6%), immune deficiency disease (IDD; 1.3 and 4.8%), and Cardiovascular disease (9.1 and 28.9%), each. Finally, the mean of hospitalization duration was 6.35 days (SD = 5.87) for patients under 60 and 8.8 days (SD = 5.9) for patients over 60.

Measurements

General Health Questionnaire

The self-report GHQ-12 was developed to screen global mental state (Goldberg et al., 1997). Among two common scoring methods of the bi-modal (0–0–1–1) and Likert scoring (0–1–2–3) types, the Likert method is preferable since it measures the symptom severity on a continuum (Hystad and Johnsen, 2020). In this study, the scoring based on the 4-point Likert-scale (0–1–2–3) was used, in which: 0 = “not at all,” 1 = “no more than usual,” 2 = “rather more than usual,” and 3 = “much more than usual,” where a higher score indicated lower mental health (Goldberg et al., 1997).

Pittsburgh Sleep Quality Index

This 19-item self-administered tool (Buysse et al., 1989) was designed for brief assessment of seven components: (1) subjective sleep quality (e.g., “how would you rate your sleep quality overall?”), (2) sleep latency (e.g., “cannot get to sleep within 30 min”), (3) sleep duration (e.g., “how many hours of actual sleep do you get at night?”), (4) sleep efficiency (e.g., “when have you usually gone to bed?”), (5) sleep disturbances (e.g., “wake up in the middle of the night or early morning”), (6) use of sleeping medication (e.g., “how often have you taken medicine to help you sleep?”), and (7) daytime dysfunction (e.g., “how often have you had trouble staying awake while driving...”). Each component was weighted on a Likert scale from 0 to 3, with higher scores indicating poorer sleep quality. Cronbach's alpha in the current study was .77 for the total score. The Persian version of PSQI that showed adequate psychometric properties (Farrahi Moghaddam et al., 2012) was used in the current study.

Perceived Stress Scale

PSS is a 10-item unidimensional scale (Cohen et al., 1983) that measures how much patients appraise the situations in their life as stressful during the preceding month. Items were coded based on a 5-point Likert-type scale: 0 (never), 1 (almost never), 2 (once in a while), 3 (often), and 4 (very often). Higher scores indicated higher perceived stress (e.g., “unable to control the important things in your life”). Cronbach's α for the Persian version of PSS was .84 for the total

score (Maroufizadeh et al., 2018), while an alpha value of .68 was obtained in our study.

ADL-Katz Scale and The Lawton Instrumental Activities of Daily Living Scale

The 6-item ADL-Katz Scale (Katz et al., 1970) assessed the ability of bathing, transferring, dressing and grooming, walking, toileting, and feeding in people over 60 years of age. The Lawton Instrumental Activities of Daily Living Scale (Lawton, 2000) measured instrumental functioning, namely, using the phone, doing housework, doing laundry, managing transportation, shopping, cooking, managing medications, and managing finances. Items for both ADL and IADL were scored based on 0 (no) and 1 (yes). Cronbach's alphas for the Persian versions of ADL and IADL were 0.80 (Sharifi et al., 2018) and between 0.72 and 0.76 (Mehraban et al., 2014), respectively. In this study, alphas were 0.66 and 0.82 for ADL and IADL, respectively.

Medical conditions and demographic characteristics

In order to evaluate the patients' demographics and medical characteristics, a questionnaire constructed by researchers was used. The medical features included: (a) the history of underlying diseases (Diabetes, HTN, HLP, Myocardial Infarction (MI), Cerebrovascular Accident (CVA), Pulmonary Disease (PD), Kidney failure, Psychiatry disorders, Obesity, IDD, and CVD) (yes/no), (b) cigarette, alcohol, and drug history (yes/no), (c) hospitalization duration (days), (d) use of downer or sleeping pills (yes/no), and (e) the change in sleep time before and after COVID-19 (hours). The demographic characteristics included: (a) gender (male/female), (b) age (years), (c) job status (employed/unemployed), (d) educational level (illiterate, primary education, secondary education, diploma level, and higher education), and (e) the number of family members.

Procedure

The current cross-sectional study was carried out on patients hospitalized due to Coronavirus infection at the Baharloo and Ziaeeian Hospitals from March to October 2020 in Tehran, Iran. After being discharged from the hospital, those who accepted to take part in the current study were asked to sign the consent form. The demographic information of those who consented to take part and their contact number was collected in a registration form. Then, three psychologists collected the data (demographic and medical variables, GHQ-12, PSS-10, PSQI, ADL and IADL), using telephone survey. Patients were informed about their optional participation in the current research and that they can leave the research any time they wish. This study received ethic approval from the Review Board of Tehran University of Medical Sciences (Ethical Code: IR.TUMS.VCR.REC.1399.156).

Statistical strategy

Data screening was performed via IBM SPSS Statistics (Version 28). CFA tests of the GHQ-12 were conducted using Mplus version 8.8. Evaluating the assumption of normality revealed a mostly positive but non-substantial skewness in all items; thus, transformation was

not required (Gravetter et al., 2020). We applied CFA using the Weighted Least Square Mean and Variance Adjusted (WLSMV) estimator. Statistical strategies were as follows: First, we used the following statistical tests and indices (MacCallum et al., 1996; Hu and Bentler, 1999; Hooper and Coughlan, 2008) to assess the models' "goodness-of-fit" (acceptable values in parenthesis): the Chi-square (χ^2 ; desired $p > 0.05$), the Comparative Fit Index (CFI > 0.95), the Tucker–Lewis Index (TLI > 0.95), the Standardized Root Mean Square Residual (SRMR < 0.08), the Normal Chi-square ($\chi^2/df < 5$), the Root Mean Square Error of Approximation (RMSEA < 0.10), and its 90% Confidence Interval (Bentler and Bonett, 1980; MacCallum et al., 1996; Loehlin, 2004; Miles and Shevlin, 2007). The exact fit is defensible when the Chi-square is not significant, regardless of the SRMR value. Approximate fit is tenable when Chi-square is significant, SRMR ≤ 0.08 , and standard residuals are all small ($|r_{res}| < 0.1$), and finally poor fit is concluded if Chi-square is significant, and SRMR > 0.08 (Satorra and Bentler, 2010).

Second, for internal consistency—as recommended for ordinal Likert-type scales, the equivalents of Cronbach's alpha coefficient (Ordinal Theta and Omega reliability coefficients) using R version 4.1.2 (Team, R. C, 2013; Revelle, 2017) were conducted, which instead of the Pearson correlation matrix, apply the poly-choric correlation matrix (Zumbo et al., 2007; Gadermann et al., 2012). A reliability coefficient of 0.70 or higher was considered an acceptable level (Cicchetti, 1994).

Third, the criterion validity was evaluated by the Spearman coefficient of rank correlation of GHQ-12 with PSQI, PSS-10, and ADL/IADL, since the data showed evidence of non-normality. Correlation coefficients were interpreted based on the effect size classification of Cohen (1988): 0.10 = small, 0.30 = medium, 0.50 = large, and 0.70 = very large.

Forth, Multivariate Analysis of Variance (MANOVA) and effect size (Hedge's g) were used to compare the mean and standard

deviation of the GHQ-12 scores across gender. According to a rule of thumb suggested by Cohen (1988), effect sizes were classified into small (< 0.20), medium (0.21–0.50), large (0.51–0.80), and very large (> 0.80).

Results

Aim 1: GHQ-12 construct validity

To test the GHQ-12 factor structure, CFA was conducted and the goodness of fit for 13 models was examined (Table 2). Model 1 (M_1) examined a general factor, in which, the total of the 12 items were loaded on a single common factor of general mental health (Goldberg et al., 1997; Romppel et al., 2013; Gnams and Staufenbiel, 2018; Hystad and Johnsen, 2020; Alaminos-Torres et al., 2021) to test the unidimensional model of assumed latent factor and included just random measurement error and indicator-specific variance (Gustafsson and Åberg-Bengtsson, 2010). If the general factor model fitted the data well, it meant that the assumption of the multidimensionality of the measurement tool was violated. Models two to seven (M_2 to M_7) consisted of a the first-order two-factor oblique models that suggested two subscales measuring two distinct dimensions (Politi et al., 1994; Schnritz et al., 1999; Montazeri et al., 2003; Kalliath et al., 2004; Najarkolaei et al., 2014; Zhong et al., 2021; Hamad, 2022). Models 8–13 (M_8 to M_{13}) examined first-order three-factor oblique models, resembling the Exploratory Factor Analysis (EFA) according to the literature (Graetz, 1991; Farrell, 1998; Martin, 1999; Daradkeh et al., 2001; Gao et al., 2004; del Pilar Sánchez-López and Dresch, 2008; Liang et al., 2016; Lee and Kim, 2020). Model 8 (M_8) included general dysphoria, lack of enjoyment, and social dysfunction (Daradkeh et al., 2001). Model 9 (M_9) consisted of anxiety, depression,

TABLE 2 Fit indices of the Measurement Models of the GHQ-12.

Model	χ^2	df	χ^2/df	CFI	TLI	RMSEA	SRMR	Based Model	$\Delta\chi^2$ (df)
M_1 (Goldberg et al., 1997; Romppel et al., 2013; Gnams and Staufenbiel, 2018; Hystad and Johnsen, 2020; Alaminos-Torres et al., 2021)	1030.608	54	19.08	0.900	0.878	0.240 (0.227–0.253)	0.172	–	–
M_2 (Politi et al., 1994) [†]	1016.411	53	19.17	0.902	0.878	0.241 (0.228–0.254)	0.170	M_1	–
M_3 (Kalliath et al., 2004) [†]	458.688	19	24.14	0.869	0.806	0.271 (0.250–0.293)	0.099	M_1	–
M_4 (Montazeri et al., 2003) [†]	955.448	52	18.37	0.908	0.883	0.235 (0.222–0.248)	0.165	M_1	–
M_5 (Najarkolaei et al., 2014; Zhong et al., 2021) [†]	1023.869	53	19.31	0.901	0.877	0.242 (0.229–0.255)	0.170	M_1	–
M_6 (Schnritz et al., 1999) [†]	951.700	43	22.13	0.825	0.776	0.259 (0.245–0.274)	0.134	M_1	–
M_7 (Hamad, 2022) [†]	1017.709	53	19.20	0.902	0.878	0.241 (0.228–0.254)	0.172	M_1	–
M_8 (Daradkeh et al., 2001)	672.991	51	13.19	0.938	0.920	0.197 (0.184–0.210)	0.099	M_1	357.62 (3)***
M_9 (Farrell, 1998) [†]	1016.582	51	19.93	0.904	0.876	0.246 (0.233–0.259)	0.135	M_1	–
M_{10} (Graetz, 1991; Gao et al., 2004; Lee and Kim, 2020)	530.411	51	10.40	0.952	0.938	0.173 (0.160–0.187)	0.087	M_1	500.20 (3)***
M_{11} (Martin, 1999)	751.738	51	14.73	0.930	0.910	0.209 (0.196–0.223)	0.107	M_1	278.87 (3)***
M_{12} (Liang et al., 2016) [†]	3018.809	50	60.37	0.704	0.610	0.435 (0.422–0.448)	0.287	M_1	–
M_{13} (del Pilar Sánchez-López and Dresch, 2008) ^{††}	502.118	50	10.04	0.955	0.941	0.170 (0.156–0.183)	0.080	M_1	528.49 (4)***

χ^2 , Chi-square; df, degrees of freedom; TLI, Tucker–Lewis index; CFI, comparative fit index; ABIC, sample-size adjusted Bayesian information criterion; χ^2/df , normal Chi-square; $\Delta\chi^2$, difference between minus twice log likelihoods between the full and the nested models; SRMR, standardized root mean square residual; RMSEA, root mean square error of approximation; Δ , differences between parameters of two models; [†] = The problem with the model is that the factors correlate greater than one which makes the model inadmissible, then, model cannot be used. ^{††} = The final selected model. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. The bold values correspond to the final selected model.

and social dysfunction (Farrell, 1998). Model 10 (M_{10}) was loaded by all three first-order factors, which included social dysfunction, anxiety and depression, and loss of confidence (Graetz, 1991; Gao et al., 2004; Lee and Kim, 2020). Model 11 (M_{11}) included cope, stress, and low self-esteem (Martin, 1999). Model 12 (M_{12}) comprised low level of social function, anxiety/depression, and poor self-confidence (Liang et al., 2016). Finally, Model 13 (M_{13}) included successful coping, self-esteem, and stress (del Pilar Sánchez-López and Dresch, 2008).

Model selection

In Table 2, all two-factor models (Politi et al., 1994; Schnitz et al., 1999; Montazeri et al., 2003; Kalliath et al., 2004; Najarkolaei et al., 2014; Zhong et al., 2021; Hamad, 2022) are inadmissible, due to factors correlate greater than 1.00 between two latent factors. The correlation between the two latent factors in two out of six models for the three-factor model also exceeded 1.00, as shown in Table 2 (M_9 and M_{12} ; Farrell, 1998; Liang et al., 2016). The fit indices of the three-factor oblique model for remaining four models (Table 2; M_8 , M_{10} , M_{11} , and M_{13}) met some of the specified fit criteria, as prior, and based on the theory-derived models (Graetz, 1991; Martin, 1999; Daradkeh et al., 2001; Gao et al., 2004; del Pilar Sánchez-López and Dresch, 2008; Lee and Kim, 2020). Then, the parsimonious principle (Bollen, 1989) was used to compare the fit indices of the three-factor first-order oblique models (Table 2; M_8 [$\Delta\chi^2 = 357.62$, $\Delta df = 3$, $p < 0.001$], M_{10} [$\Delta\chi^2 = 500.20$, $\Delta df = 3$, $p < 0.001$], M_{11} [$\Delta\chi^2 = 278.87$, $\Delta df = 7$, $p < 0.001$], and M_{13} [$\Delta\chi^2 = 528.49$, $\Delta df = 4$, $p < 0.001$]) with those of the unidimensional first-order model (M_1) as the baseline/null model. For three-factor models, four out of six models exhibited similar fitness, though with poor goodness-of-fit (Table 2). To determine the most efficient parsimonious model, the nesting and equivalence testing (NET) methodology was implemented via Mplus 8.8 (Bentler and Satorra, 2010; Asparouhov and Muthén, 2019). Since all models are non-tested and/or non-equivalent, as was expected (the NET value = 0.0000001), it can be concluded that model 13 showed the best fit, due to its low Chi-square value in comparison with the others ($\chi^2/df = 10.04$; CFI = 0.96; TLI = 0.94; RMSEA = 0.17; 90% CI = 0.16 to 0.18; SRMR = 0.08). More information on confirmatory factor analysis of models 1, 8, 10, 11, and 13 is presented in supporting information file (Supplementary Figures S1–S5).

Aim 2: GHQ-12 reliability

The Ordinal *Theta* and the *Omega* reliability coefficients for the subscales of GHQ-12 are presented in Table 3. The means of inter-item correlation were 0.10, 0.53, 0.48, and 0.58 for the total score, successful coping, self-esteem, and stress, respectively. Almost all

items within the three subscales had a moderate positive relationship with each other (based on the corrected item-total correlation for subscale's items), with values ranging from 0.34 to 0.83, 0.23 to 0.78, and 0.53 to 0.66 for successful coping, self-esteem, and stress, respectively.

Aim 3: GHQ-12 and related measures

Table 4 demonstrates that the inter-correlation between GHQ-12 total score and subscales ranged from 0.40 to 0.83 ($p < 0.01$). Criterion validity was estimated by the correlation of GHQ-12 total score and its subscales with PSQI and PSS (Table 4). PSQI had significant positive correlations with total GHQ-12 ($r = 0.28$, $p < 0.01$), successful coping ($r = 0.24$, $p < 0.01$), self-esteem ($r = 0.21$, $p < 0.01$), and stress ($r = 0.20$, $p < 0.01$). Also, PSS had significant positive correlations with the total score of GHQ-12 and its subscales ($r = 0.31$ to 0.58 , $p < 0.01$). In the over 60 years of age group, negative correlations of the total GHQ-12 score were found with the ADL ($r = -0.34$, $p < 0.01$; $r = -0.37$, $p < 0.01$) and IADL scores ($r = -0.42$, $p < 0.01$, $r = -0.46$, $p < 0.01$) before and after the infection of COVID-19, respectively (Table 4).

As depicted in Table 5 for demographic and medical variables, the total score of GHQ-12 was significantly correlated with HLP ($r = 0.16$, $p < 0.01$), kidney failure ($r = 0.12$, $p < 0.05$), psychiatry disorders ($r = 0.18$, $p < 0.01$), the hospitalization duration ($r = 0.15$, $p < 0.01$), the change in sleep time ($r = 0.31$, $p < 0.01$), use of sleeping pills ($r = 0.30$, $p < 0.01$), educational level ($r = -0.26$, $p < 0.01$), and the number of family members ($r = -0.12$, $p < 0.05$). The correlation of the total score of GHQ-12 with Diabetes, HTN, MI, CVA, PD, obesity, IDD, CVD, and drug, alcohol, and cigarette history were non-significant ($p > 0.01$). Information for the correlation of GHQ-12 subscales is presented in Table 5.

Aim 4: Gender, age, and GHQ-12

Table 3 presents the mean and Standard Deviations (SD) of the GHQ-12 total score and subscales across gender and age groups. The female patients scored significantly higher than the males on total GHQ-12 scores [$t(310) = -4.77$, $p < 0.001$, Cohen's $d = -0.65$, mean difference bootstrap 95% CI = -4.46 to -1.92]. The patients above 60 also scored slightly higher than patients under 60 on their total GHQ-12 scores [$t(312) = -1.45$, $p = 0.08$, Cohen's $d = -0.19$, mean difference bootstrap 95% CI = -2.94 to 0.55]. A Multivariate Analysis of Variance (MANOVA) showed significant group differences by gender [$F(3-308) = 10.80$, $p < 0.001$, $\eta^2 = 0.095$] and age groups

TABLE 3 The Descriptive statistics of GHQ-12.

	Mean (SD)					Ordinal Theta	Omega	α
	Total	Female	Male	Under 60	Over 60			
1. Successful Coping	8.95 (3.42)	8.39 (3.20)	9.39 (3.55)	9.30 (3.31)	7.98 (3.55)	0.90	0.86	0.87
2. Self-esteem	3.91 (2.63)	4.37 (2.61)	3.22 (2.39)	3.86 (2.54)	4.04 (2.88)	0.75	0.77	0.77
3. Stress	3.96 (2.43)	4.72 (2.40)	3.33 (2.27)	4.06 (2.40)	3.67 (2.51)	0.93	0.88	0.80
4. GHQ-12	15.52 (6.19)	16.18 (3.66)	14.87 (3.82)	15.88 (3.80)	14.39 (3.82)	-	-	0.86

Notes. GHQ = General health questionnaire, α = Alpha, SD = Standard deviation. * $p < 0.05$, ** $p < 0.01$.

TABLE 4 The correlation between GHQ-12 subscales and their correlations with sleep quality, perceived stress, and daily functioning.

	2	3	4	PSQI						PSS		ADL		IADL		
				Total	SSQ	SL	SD	SE	SD	USM	DD	PSS	Before	After	Before	After
1. Successful Coping	0.40**	0.45**	0.83**	0.24**	0.15	0.26**	0.29**	0.25**	0.51**	0.31**	-0.35**	-0.43**	-0.48**	-0.55**		
2. Self-esteem		0.73**	0.80**	0.27**	0.20*	0.22*	0.27**	0.23**	0.23**	0.50**	-0.25*	-0.33*	-0.28**	-0.31**		
3. Stress			0.81**	0.23**	0.38*	0.27**	0.48**	0.23**	0.28**	0.58**	-0.22*	-0.16	-0.19	-0.15		
4. GHQ-12				0.30**	0.28**	0.30**	0.42**	0.30**	0.45**	0.53**	-0.34**	-0.37**	-0.42**	-0.46**		

Notes: GHQ = General health questionnaire, PSQI = Pittsburgh sleep quality index, SSQ = Subjective sleep quality, SL = Sleep latency, SD = Sleep duration, SE = Sleep efficiency, USM = Use of sleeping medication, DD = Daytime dysfunction, PSS = Perceived stress scale. * $p < 0.05$, ** $p < 0.01$.

[$F(3-310) = 7.27, p < 0.001, \eta^2 = 0.066$] on mean scores of the subscales (see Table 3 for Mean scores). Subsequent tests of between-subjects' effects showed that females scored significantly higher on successful coping [$F(1-310) = 5.68, p < 0.05, \eta^2 = 0.018$], self-esteem [$F(1-310) = 28.11, p < 0.001, \eta^2 = 0.083$], and stress [$F(1-310) = 27.38, p < 0.081, \eta^2 = 0.081$] than males. The patients over 60 scored significantly higher on successful coping [$F(1-312) = 9.41, p < 0.01, ns, \eta^2 = 0.029$], than patients under 60. However, non-significant differences in mean score were found for self-esteem [$F(1-312) = 0.31, p = 0.58, ns, \eta^2 = 0.001$] and stress [$F(1-312) = 1.60, p = 0.21, ns, \eta^2 = 0.005$] across age.

Discussion

The present study sought to evaluate the psychometric properties of the General Health Questionnaire-12 in patients hospitalized with the diagnosis of COVID-19. Overall, our results provided support for the construct and criterion validity, internal consistency, and invariance of GHQ-12 across age (under 60 vs. over 60) and gender (male vs. female). Therefore, this questionnaire showed that it can be applied in Iranian COVID-19 patients.

Among 13 theoretically and empirically emerged models of the GHQ-12 tested in this study, the current data fitted better with the three-factor model of del Pilar Sánchez-López and Dresch (2008), including successful coping, self-esteem, and stress. The factor loading of all items was adequate (Ford et al., 1986). This result is contrary to Liang et al. (2016) study that showed equal model fit (CFI=0.98, RMSEA=0.03) for 11 previously emerged factorial models. The unidimensional model (Goldberg et al., 1997) was not supported in our study, suggesting that GHQ-12 may not be a homogeneous tool that measures only one construct of mental health, or rather, it may cover several constructs instead of concentrating on “narrow aspects” of mental health (Gustafsson and Åberg-Bengtsson, 2010). The established factorial model manifested good reliabilities. Indeed, the overall results of the Cronbach's alpha, Theta, and Omega coefficients were satisfactory, with the adequate means of inter-item correlations for subscales. These results are in line with a bulk of studies on the psychometric features of GHQ-12 in different contexts (Liang et al., 2016; Elovania et al., 2020).

To test how accurately the GHQ-12 can correlate the expected outcomes, the criterion validity was conducted as our third objective through the relationship of GHQ with perceived stress, sleep quality, ADL/IADL, and demographic and medical variables. First, GHQ-12 total score and subscales showed significant weak to strong correlations with the sleep quality total score and all subscales, where the subscale of subjective sleep quality had the strongest correlation coefficients. These findings are supported by previous research (Xiong et al., 2019; Aquil et al., 2021; Thielmann et al., 2021). Oh et al. (2019), for instance, found that adults with higher psychological distress had higher difficulty falling asleep. Second, perceived stress showed moderate to high positive correlations with the GHQ-12 total score and three subscales, further supporting the criterion validity of GHQ-12. These findings were consistent with previous studies (Örücü and Demir, 2009; Gajula et al., 2021). It is thought that psychosocial stressors, such as living alone, social restrictions and isolation, financial burden, and loss of family members, that lead to heightened anxiety, fear, and

TABLE 5 The correlations of Mental Health with demographic and medical variables.

	HLP	KF	Psychiatry disorders	Hospitalization duration	Change in sleep time	Use of sleeping pills	Educational level	Family members (n)
1. Successful Coping	0.14*	0.09	0.13*	0.17**	0.10	0.24**	-0.30**	-0.12*
2. Self-esteem	0.15**	0.08	0.14*	0.10	0.29**	0.25**	-0.17**	-0.11*
3. Stress	0.06	0.09	0.17**	0.08	0.44**	0.25**	-0.08	-0.05
4. GHQ-12	0.16**	0.12*	0.18**	0.15**	0.31**	0.30**	-0.26**	-0.12*

Notes: GHQ = General health questionnaire, HLP = Hyperlipidemia, KF = Kidney failure. * $p < 0.05$, ** $p < 0.01$.

anger, made a significant contribution to the higher level of stress experienced by patients with COVID-19 (Torales et al., 2020; Matalon et al., 2021; Varman et al., 2022).

The GHQ-12 in patients over 60 demonstrated negative correlations with ADL and IADL-as the third criterion variable. A similar result was found by earlier studies (Albanese et al., 2020). Our finding suggested that lack of autonomy in daily life can seriously damage a person's self-esteem, increase their conflicts with others, and make them more vulnerable to symptoms of anxiety and depression.

The fourth variable used to evaluate the criterion validity of GHQ-12 was medical variables, that manifested a set of significant associations. This questionnaire had significant positive correlation with HLP, in line with the study of Wang et al. (2016) that found the GHQ-12 scores was significantly higher in patients with HLP. This association seems bidirectional, given that on the one hand, the empirical evidence suggested that HLP triggers the onset of depression (Chuang et al., 2014), and on the other hand, patients with depression experience a higher incidence of HLP, compared to the general population (Chien et al., 2013). Chang et al. (2021) also showed positive correlation of high blood fat and stress. Kidney failure was shown to be positively correlated with GHQ-12, consistent with two systematic reviews that found a high rate of depression in patients with Kidney failure (Bautovich et al., 2014; Kondo et al., 2020). One explanation for this link might be the impact of difficulties the patients with kidney failure face, such as the psychological and social burden of the disease, comorbid diseases, and the experience of dialysis, which may lead to depression/anxiety (Ozcan et al., 2015). These symptoms are, in turn, associated with negative outcomes including poor quality of life, poor treatment compliance, and elevated mortality rates (Bautovich et al., 2014; Butt et al., 2022). Furthermore, GHQ-12 was significantly correlated with psychiatry disorders. This association is expected because mental illnesses decrease the quality of life and severely impair patients' ability to communicate and form social relationships. Therefore, it is likely that they are more affected by a pandemic than those with no psychiatric conditions (Kaufman et al., 2020). GHQ-12 was positively correlated with the hospitalization duration. This is another expected result, because at hospitals, patients experience a loss of dignity as a result of their physical conditions, elevating their senses of powerlessness, embarrassment, and being violated. Consequently, these debilitating experiences may lead to mental distress (Liao et al., 2020). Finally, GHQ-12 showed positive correlations with change in sleep time and use of sleeping pills. The change in the sleep-wake cycle might be explained by a third mechanism like anxiety symptoms (Becker et al., 2018). Tang et al. (2017) suggested that higher score of GHQ-12 was strongly predicted by the reduced sleep duration and use of sleeping pills, and vice versa. They argued that extremely long or short sleep duration and excessive use of sleeping pills lead to difficulty in daytime function, which result in adverse outcomes. However, it should be noted that all of these correlations in the current study were weak to moderate and should be interpreted cautiously.

Finally, the criterion validity of GHQ-12 was examined via its link with demographic characteristics. GHQ-12 showed negative correlations with educational level and the number of family members. Consistent with our finding, Dalgard et al. (2007) study have shown a

significant association between lower educational level and psychological distress in both Norwegian males and female. Additionally, people who live alone may be especially dependent on others for social connection and support, making them more vulnerable to social distancing (Hendriksen et al., 2021). Hence, larger number of family members may be a protective factor against the sense of loneliness and act as a means for social support.

Due to some clues that showed the gender and age differences in the level of GHQ-12, we investigated these group mean differences to be considered in the future use of the questionnaire. In the present study, consistent with previous evidence (Giorgi et al., 2014), lower average of general health in women (higher GHQ-12 total scores and subscales) was observed. Previous evidence have shown that the prevalence of factors thought to be intensified during a pandemic (such as preceding anxiety and depression, chronic environmental exposure, and domestic violence) is higher among women (Bucciarelli et al., 2021). This could increase women's odds of developing mental health issues. As for age differences, higher level of the GHQ-12 total score and successful coping subscale in patients older than 60 years was found in the present study. This result contradicted previous findings that indicated the association of aging with an intrinsic reduction in susceptibility to psychological distress (Hoeymans et al., 2004). However, losing social contacts in the elderly (e.g., the death of family members), becoming prohibited from engagement in common social interactions due to social distancing order, and receiving limited access to social support and services may increase their susceptibility to mental health problems (Stuart et al., 2022).

Limitations, future directions, and clinical implications

The current study results provide insight into the general mental health status in patients with COVID-19. However, this study is not without limitations. First, GHQ-12 is a screening tool and was not designed for diagnosis objectives and distinguishing among mental disorders (Goldberg, 1986; Schnitz et al., 1999). Researchers in future work can use semi-structured interviews to provide more in-depth information regarding high scored items of GHQ-12 and compare yielded scores of GHQ-12 with the additional probe questions (i.e., regarding symptom severity and duration) to evaluate the accuracy of GHQ-12. Second, the cross-sectional design of the current study has prevented causal inferences. It restricts our knowledge on the direction of the correlation of GHQ-12 with perceived stress and sleep quality. It also prevents us from measuring the stability of mental health scores over time. Longitudinal studies are recommended to explore the predictive role of the abovementioned variables on each other, and evaluate the stability of GHQ-12 scores over time. Finally, the present study did not perform the measurement invariance analyses across gender or age, because the sample sizes would start to become small when we divided the sample into age and gender subgroups. In the age case, sample size would be down to 87 for the younger segment. Such small sample sizes lacked sufficient power to detect any invariance.

Given the positive link of poor mental health with perceived stress, sleep disturbances, and impaired independent daily activities among Iranian COVID-19 patients, an important clinical implication for physicians, psychologists, and psychiatrists is to

design psychological interventional courses for COVID-19 hospitalized patients that specifically target these problems. Access to such services *via* social media may be beneficial not only for their mental health, but also for their ability to improve their physical and mental functioning and independency (Shojaei and Masoumi, 2020).

Conclusion

The current study was undertaken to evaluate the psychometric properties of GHQ-12 among Iranian COVID-19 patients. The results of construct validity analyses supported the three-factor model of successful coping, self-esteem, and stress, which showed satisfactory reliability. The criterion validity of GHQ-12 was confirmed through its positive relationship with perceived stress and sleep quality, as well as its negative relationships with activities and instrumental activities of daily living in patients with over 60 years of age. Women and patients above 60 manifested higher GHQ-12 scores, compared to men and patients under 60.

Data availability statement

The raw data supporting the conclusions of this article can be provided by the corresponding author upon reasonable request.

Ethics statement

The studies involving human participants were reviewed and approved by the Review Board of Tehran University of Medical Sciences (ethical code: IR.TUMS.VCR.REC.1399.156). The patients/participants provided their written informed consent to participate in this study.

Author contributions

MHA, FE, and ZV: conceptualization, design, methodology, and investigation and project administration. FF and FG: data collection. MHA: formal analysis and supervision. PSY, NAS, and MHA: writing the original draft. PSY, NW, and MHA: revising the draft. All authors have contributed to the conception and design of the study, drafted or revised this manuscript, reviewed the final version of this manuscript before submission, and agreed to be accountable for all aspects of the work.

Funding

This research was supported by grant number 99-1-101-47,395 from the Tehran University of Medical sciences.

Acknowledgments

The authors would like to thank all patients who participated in the study and the Baharloo and Ziaiean Hospitals' staff who assisted in data collection.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations,

or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpsyg.2023.1132154/full#supplementary-material>

References

- Achdut, N., and Refaeli, T. (2020). Unemployment and psychological distress among young people during the COVID-19 pandemic: psychological resources and risk factors. *Int. J. Environ. Res. Public Health* 17:7163. doi: 10.3390/ijerph17197163
- Alaminos-Torres, A., Martínez-Lorca, M., Sifre De Sola, I., López-Ejeda, N., and Dolores Marrodán, M. (2021). Psychological distress in Spanish airline pilots during the aviation crisis caused by the COVID-19 pandemic and psychometric analysis of the 12-item general health questionnaire. *Ergonomics* 65, 741–752. doi: 10.1080/00140139.2021.1984587
- Albanese, A. M., Bartz-Overman, C., Parikh, M. T., and Thielke, S. M. (2020). Associations between activities of daily living independence and mental health status among medicare managed care patients. *J. Am. Geriatr. Soc.* 68, 1301–1306. doi: 10.1111/jgs.16423
- Aquil, A., El Kherchi, O., El Azmaoui, N., Mouallif, M., Guerroumi, M., Chokri, A., et al. (2021). Body image dissatisfaction and lower self-esteem as major predictors of poor sleep quality in gynecological cancer patients after surgery: cross-sectional study. *BMC Womens Health* 21, 229–231. doi: 10.1186/s12905-021-01375-5
- Asparouhov, T., and Muthén, B. (2019). Nesting and equivalence testing for structural equation models. *Struct. Equ. Model. Multidiscip. J.* 26, 302–309. doi: 10.1080/10705511.2018.1513795
- Bautovich, A., Katz, I., Smith, M., Loo, C. K., and Harvey, S. B. (2014). Depression and chronic kidney disease: a review for clinicians. *Aust N Z J Psychiatry* 48, 530–541. doi: 10.1177/0004867414528589
- Becker, S. P., Jarrett, M. A., Luebbe, A. M., Garner, A. A., Burns, G. L., and Kofler, M. J. (2018). Sleep in a large, multi-university sample of college students: sleep problem prevalence, sex differences, and mental health correlates. *Sleep Health* 4, 174–181. doi: 10.1016/j.sleh.2018.01.001
- Bentler, P. M., and Bonnett, D. G. (1980). Significance tests and goodness of fit in the analysis of covariance structures. *Psychol. Bull.* 88, 588–606. doi: 10.1037/0033-2909.88.3.588
- Bentler, P. M., and Satorra, A. (2010). Testing model nesting and equivalence. *Psychol. Methods* 15, 111–123. doi: 10.1037/a0019625
- Bollen, K. A. (1989). *Structural Equations with Latent Variables*. New York: John Wiley & Sons.
- Bonsaksen, T., Leung, J., Price, D., Ruffolo, M., Lamph, G., Kabelenga, I., et al. (2022). Self-reported Long COVID in the general population: sociodemographic and health correlates in a cross-national sample. *Life* 12:901. doi: 10.3390/life12060901
- Bruine de Bruin, W. (2021). Age differences in COVID-19 risk perceptions and mental health: evidence from a national US survey conducted in March 2020. *J Gerontol Ser B* 76, e24–e29. doi: 10.1093/geronb/gbaa074
- Bucciarelli, V., Nasi, M., Bianco, F., Seferovic, J., Ivkovic, V., Gallina, S., et al. (2021). Depression pandemic and cardiovascular risk in the COVID-19 era and long COVID syndrome: gender makes a difference. *Trends Cardiovasc. Med.* 32, 12–17. doi: 10.1016/j.tcm.2021.09.009
- Butt, M. D., Ong, S. C., Butt, F. Z., Sajjad, A., Rasool, M. F., Imran, I., et al. (2022). Assessment of health-related quality of life, medication adherence, and prevalence of depression in kidney failure patients. *Int. J. Environ. Res. Public Health* 19:15266. doi: 10.3390/ijerph192215266
- Byusse, D. J., Reynolds, C. F. III, Monk, T. H., Berman, S. R., and Kupfer, D. J. (1989). The Pittsburgh sleep quality index: a new instrument for psychiatric practice and research. *Psychiatry Res.* 28, 193–213. doi: 10.1016/0165-1781(89)90047-4
- Carfi, A., Bernabei, R., and Landi, F. (2020). Persistent symptoms in patients after acute COVID-19. *JAMA* 324, 603–605. doi: 10.1001/jama.2020.12603
- Chang, P.-Y., Chiou, S.-T., Lo, W.-Y., Huang, N., and Chien, L.-Y. (2021). Stressors and level of stress among different nursing positions and the associations with hyperlipidemia, hyperglycemia, and hypertension: a national questionnaire survey. *BMC Nurs.* 20, 250–210. doi: 10.1186/s12912-021-00777-y
- Chien, I.-C., Lin, C.-H., Chou, Y.-J., and Chou, P. (2013). Increased risk of hyperlipidemia in patients with major depressive disorder: a population-based study. *J. Psychosom. Res.* 75, 270–274. doi: 10.1016/j.jpsychores.2013.06.003
- Chuang, C.-S., Yang, T.-Y., Muo, C.-H., Su, H.-L., Sung, F.-C., and Kao, C.-H. (2014). Hyperlipidemia, statin use and the risk of developing depression: a nationwide retrospective cohort study. *Gen. Hosp. Psychiatry* 36, 497–501. doi: 10.1016/j.genhosppsych.2014.05.000
- Cicchetti, D. V. (1994). Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instruments in psychology. *Psychol. Assess.* 6, 284–290. doi: 10.1037/1040-3590.6.4.284
- Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences*. Abingdon-on-Thames: Routledge Academic.
- Cohen, S., Kamarck, T., and Mermelstein, R. (1983). A global measure of perceived stress. *J. Health Soc. Behav.* 24:385. doi: 10.2307/2136404
- Crayne, M. P. (2020). The traumatic impact of job loss and job search in the aftermath of COVID-19. *Psychol. Trauma Theory Res. Pract. Policy* 12, S180–S182. doi: 10.1037/tra0000852
- Czeisler, M. É., Lane, R. I., Petrosky, E., Wiley, J. F., Christensen, A., Njai, R., et al. (2020). Mental health, substance use, and suicidal ideation during the COVID-19 pandemic—United States, June 24–30, 2020. *Morb. Mortal. Wkly Rep.* 69, 1049–1057. doi: 10.15585/mmwr.mm6932a1
- Dalgard, O. S., Mykletun, A., Rognerud, M., Johansen, R., and Zahl, P. H. (2007). Education, sense of mastery and mental health: results from a nation wide health monitoring study in Norway. *BMC Psychiatry* 7, 1–9. doi: 10.1186/1471-244x-7-20
- Daradkeh, T. K., Ghubash, R., and El-Rufaie, O. E. (2001). Reliability, validity, and factor structure of the Arabic version of the 12-item general health questionnaire. *Psychol. Rep.* 89, 85–94. doi: 10.2466/pr0.2001.89.1.85
- de Castroe Costa, E., Uchoa, E., Oliveira Araújo Firmo, J., Lima-Costa, M. F., and Prince, M. (2008). Association of cognitive impairment, activity limitation with latent traits in the GHQ-12 in the older elderly. The Bambui health and aging study (BHAS). *Aging Clin. Exp. Res.* 20, 562–568. doi: 10.1023/b:wjns.2020.104019
- de Paiva Teixeira, L. E. P., de Freitas, R. L., Abad, A., da Silva, J. A., Antonelli-Ponti, M., Bastos, S., et al. (2020). Psychological impacts related to stress and fear during the COVID-19 pandemic: cardiovascular diseases, diabetes and psychological disorders as risk factors. *World J Neurosci* 10, 191–205. doi: 10.4236/wjns.2020.104019
- del Pilar Sánchez-López, M., and Dresch, V. (2008). The 12-item general health questionnaire (GHQ-12): reliability, external validity and factor structure in the Spanish population. *Psicothema* 20, 839–843.
- Deng, J., Zhou, F., Hou, W., Silver, Z., Wong, C. Y., Chang, O., et al. (2021). The prevalence of depression, anxiety, and sleep disturbances in COVID-19 patients: a meta-analysis. *Ann. N. Y. Acad. Sci.* 1486, 90–111. doi: 10.1111/nyas.14506
- Edemekong, P.F., Bomgaars, D.L., and Levy, S.B. (2017). *Activities of daily living (ADLs)*. Treasure Island (FL): StatPearls Publishing.
- Elovainio, M., Hakulinen, C., Pulkki-Råback, L., Aalto, A.-M., Virtanen, M., Partonen, T., et al. (2020). General health questionnaire (GHQ-12), Beck depression inventory (BDI-6), and mental health index (MHI-5): psychometric and predictive properties in a Finnish population-based sample. *Psychiatry Res.* 289:112973. doi: 10.1016/j.psychres.2020.112973
- Farrahi Moghaddam, J., Nakhaee, N., Sheibani, V., Garrusi, B., and Amirakafi, A. (2012). Reliability and validity of the Persian version of the Pittsburgh sleep quality index (PSQI-P). *Sleep Breath* 16, 79–82. doi: 10.1007/s11325-010-0478-5
- Farrell, G. A. (1998). The mental health of hospital nurses in Tasmania as measured by the 12-item general health questionnaire. *J. Adv. Nurs.* 28, 707–712. doi: 10.1046/j.1365-2648.1998.00735.x

- Ford, J. K., MacCallum, R. C., and Tait, M. (1986). The application of exploratory factor analysis in applied psychology: a critical review and analysis. *Pers. Psychol.* 39, 291–314. doi: 10.1111/j.1744-6570.1986.tb00583.x
- Gademann, A. M., Guhn, M., and Zumbo, B. D. (2012). Estimating ordinal reliability for Likert-type and ordinal item response data: a conceptual, empirical, and practical guide. *Pract Assess Res Eval* 17:3. doi: 10.7275/n560-j767
- Gajula, M., Bant, D., and Bathija, G. V. (2021). Perceived stress among adolescent school students in Hubli: a cross-sectional study. *Natl J Community Med* 12, 169–174. doi: 10.5455/njcm.20210625054326
- Gao, F., Luo, N., Thumboo, J., Fones, C., Li, S.-C., and Cheung, Y.-B. (2004). Does the 12-item general health questionnaire contain multiple factors and do we need them? *Health Qual. Life Outcomes* 2, 1–7. doi: 10.1186/1477-7525-2-63
- Giorgi, G., Perez, J. M. L., D'Antonio, A. C., Perez, F. J. F., Arcangeli, G., Cupelli, V., et al. (2014). The General Health Questionnaire (GHQ-12) in a sample of Italian workers: mental health at individual and organizational level. *World J Med Sci* 11, 47–56. doi: 10.5829/idosi.wjms.2014.11.1.83295
- Gnams, T., and Staufenbiel, T. (2018). The structure of the general health questionnaire (GHQ-12): two meta-analytic factor analyses. *Health Psychol. Rev.* 12, 179–194. doi: 10.1080/17437199.2018.1426484
- Goldberg, D. P. (1972). The detection of psychiatric illness by questionnaire. *Maudsley Monogr.* Windsor, UK: Nfer-Nelson Publisher, 21:483.
- Goldberg, D. (1986). Use of the general health questionnaire in clinical work. *Br. Med. J. (Clin. Res. Ed.)* 293, 1188–1189. doi: 10.1136/bmj.293.6556.1188
- Goldberg (1988). *User's Guide to the General Health Questionnaire*. Windsor.
- Goldberg, D. P., Gater, R., Sartorius, N., Ustun, T. B., Piccinelli, M., Gureje, O., et al. (1997). The validity of two versions of the GHQ in the WHO study of mental illness in general health care. *Psychol. Med.* 27, 191–197. doi: 10.1017/s0033291796004242
- Graetz, B. (1991). Multidimensional properties of the general health questionnaire. *Soc. Psychiatry Psychiatr. Epidemiol.* 26, 132–138. doi: 10.1007/bf00782952
- Gravetter, F. J., Wallnau, L. B., Forzano, L.-A. B., and Witnauer, J. E. (2020). *Essentials of Statistics for the Behavioral Sciences*. Boston, MA: Cengage Learning.
- Gustafsson, J.-E., and Åberg-Bengtsson, L. (2010). "Unidimensionality and interpretability of psychological instruments," in *Measuring Psychological Constructs: Advances in Model-Based Approaches*. ed. S. E. Embretson (Washington, DC, US: American Psychological Association), 97–121.
- Halpin, D. M., Singh, D., and Hadfield, R. M. (2020). Inhaled corticosteroids and COVID-19: a systematic review and clinical perspective. *Eur. Respir. J.* 55:2001009. doi: 10.1183/13993003.01009-2020
- Hamad, E. O. (2022). Psychological well-being among Saudi adults during the context of COVID-19 lockdown: a psychometric analysis of the 12-item General Health Questionnaire. *BMC Psychol.* 10:319. doi: 10.21203/rs.3.rs-1591586/v1
- Hendriksen, P. A., Kiani, P., Garssen, J., Bruce, G., and Verster, J. C. (2021). Living alone or together during lockdown: association with mood, immune fitness and experiencing COVID-19 symptoms. *Psychol. Res. Behav. Manag.* 14, 1947–1957. doi: 10.2147/prbm.s332853
- Hoeymans, N., Garssen, A. A., Westert, G. P., and Verhaak, P. F. (2004). Measuring mental health of the Dutch population: a comparison of the GHQ-12 and the MHI-5. *Health Qual. Life Outcomes* 2, 1–6. doi: 10.1186/1477-7525-2-23
- Hooper, D., and Coughlan, J. (2008). Structural equation modelling: guidelines for determining model fit. *Electron. J. Bus. Res. Methods* 6, 53–60. doi: 10.21427/D7CF7R
- Hossain, M. M., Rahman, M., Trisha, N. F., Tasnim, S., Nuzhath, T., Hasan, N. T., et al. (2021). Prevalence of anxiety and depression in South Asia during COVID-19: a systematic review and meta-analysis. *Heliyon* 7:e06677. doi: 10.31234/osf.io/q4k5b
- Hu, L. T., and Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: conventional criteria versus new alternatives. *Struct. Equ. Model. Multidiscip. J.* 6, 1–55. doi: 10.1080/10705519909540118
- Hystad, S. W., and Johnsen, B. H. (2020). The dimensionality of the 12-item general health questionnaire (GHQ-12): comparisons of factor structures and invariance across samples and time. *Front. Psychol.* 11:1300. doi: 10.3389/fpsyg.2020.01300
- Kalliath, T. J., O'Driscoll, M. P., and Brough, P. (2004). A confirmatory factor analysis of the General Health Questionnaire-12. *Stress Health* 20, 11–20. doi: 10.1002/smi.993
- Katz, S., Downs, T. D., Cash, H. R., and Grotz, R. C. (1970). Progress in development of the index of ADL. *Gerontologist* 10, 20–30. doi: 10.1093/geront/10.1_part_1.20
- Kaufman, K. R., Petkova, E., Bhui, K. S., and Schulze, T. G. (2020). A global needs assessment in times of a global crisis: world psychiatry response to the COVID-19 pandemic. *BJPsych Open* 6:e48. doi: 10.1192/bjo.2020.25
- Kondo, K., Antick, J. R., Ayers, C. K., Kansagara, D., and Chopra, P. (2020). Depression screening tools for patients with kidney failure: a systematic review. *Clin. J. Am. Soc. Nephrol.* 15, 1785–1795. doi: 10.2215/CJN.05540420
- Korkut, S. (2022). Evaluation of Psychopathological Conditions, Corona Anxiety, Death Anxiety and Suicide Risk in Hemodialysis Patients during the COVID-19 Pandemic. doi: 10.21203/rs.3.rs-1211968/v1
- Lawton, M. (2000). Lawton instrumental activities of daily living scale. *Merck Manual Geriatr* 108, 52–62. doi: 10.1097/01.naj.00000314810.46029.74
- Lee, B., and Kim, Y. E. (2020). Factor structure of the 12-item general health questionnaire (GHQ-12) among Korean university students. *Psychopharmacol. Clin. Psychopharmacol* 30, 248–253. doi: 10.5455/pcp.20200518112914
- Liang, Y., Wang, L., and Yin, X. (2016). The factor structure of the 12-item general health questionnaire (GHQ-12) in young Chinese civil servants. *Health Qual. Life Outcomes* 14, 136–139. doi: 10.1186/s12955-016-0539-y
- Liao, K.-M., Ho, C.-H., Lai, C.-C., Chao, C.-M., Chiu, C.-C., Chiang, S.-R., et al. (2020). The association between depression and length of stay in the intensive care unit. *Medicine* 99:e20514. doi: 10.1097/md.00000000000020514
- Loehlin, J. C. (2004). *Latent Variable Models: An introduction to Factor, Path, and Structural Equation Analysis. 4th.* New Jersey: Lawrence Erlbaum Associates Publishers.
- MacCallum, R. C., Browne, M. W., and Sugawara, H. M. (1996). Power analysis and determination of sample size for covariance structure modeling. *Psychol. Methods* 1, 130–149. doi: 10.1037/1082-989X.1.2.130
- Maroufizadeh, S., Foroufard, F., Navid, B., Ezabadi, Z., Sobati, B., and Omani-Samani, R. (2018). The perceived stress scale (PSS-10) in women experiencing infertility: a reliability and validity study. *Middle East Fertil Soc J* 23, 456–459. doi: 10.1016/j.mefs.2018.02.003
- Maroufizadeh, S., Pourshaikhian, M., Pourramzani, A., Sheikholeslami, F., Moghadamnia, M.T., and Alavi, S.A. (2022). Prevalence of Anxiety and Depression in General Population of Iran during the COVID-19 Pandemic: A Web-Based Cross-Sectional Study. *Iran. J. Psychiatry* 17, 230–239. doi: 10.21203/rs.3.rs-39082/v1
- Martin, A. J. (1999). Assessing the multidimensionality of the 12-item general health questionnaire. *Psychol. Rep.* 84, 927–935. doi: 10.2466/pr0.1999.84.3.927
- Marvaldi, M., Mallet, J., Dubertret, C., Moro, M. R., and Guessoum, S. B. (2021). Anxiety, depression, trauma-related, and sleep disorders among healthcare workers during the COVID-19 pandemic: a systematic review and meta-analysis. *Neurosci. Biobehav. Rev.* 126, 252–264. doi: 10.1016/j.neubiorev.2021.03.024
- Matalon, N., Dorman-Ilan, S., Hasson-Ohayon, I., Hertz-Palmor, N., Shani, S., Basel, D., et al. (2021). Trajectories of post-traumatic stress symptoms, anxiety, and depression in hospitalized COVID-19 patients: a one-month follow-up. *J. Psychosom. Res.* 143:110399. doi: 10.1016/j.jpsychores.2021.110399
- Mehraban, A. H., Soltanmohamadi, Y., Akbarfahimi, M., and Taghizadeh, G. (2014). Validity and reliability of the Persian version of Lawton instrumental activities of daily living scale in patients with dementia. *Med. J. Islam Repub. Iran* 28:25.
- Miles, J., and Shevlin, M. (2007). A time and a place for incremental fit indices. *Personal. Individ. Differ.* 42, 869–874. doi: 10.1016/j.paid.2006.09.022
- Moghanibashi-Mansourieh, A. (2020). Assessing the anxiety level of Iranian general population during COVID-19 outbreak. *Asian J. Psychiatr.* 51:102076. doi: 10.1016/j.ajp.2020.102076
- Montazeri, A., Harirchi, A. M., Shariati, M., Garmaroudi, G., Ebadi, M., and Fateh, A. (2003). The 12-item general health questionnaire (GHQ-12): translation and validation study of the Iranian version. *Health Qual. Life Outcomes* 1, 1–4. doi: 10.1186/1477-7525-1-66
- Moradian, S., Teufel, M., Jahre, L., Musche, V., Fink, M., Dinse, H., et al. (2021). Mental health burden of patients with diabetes before and after the initial outbreak of COVID-19: predictors of mental health impairment. *BMC Public Health* 21, 2068–2011. doi: 10.1186/s12889-021-12101-z
- Najarkolaie, F. R., Raiisi, F., Rahnama, P., Fesharaki, M. G., Zamani, O., Jafari, M. R., et al. (2014). Factor structure of the Iranian version of 12-item general health questionnaire. *Iran Red Crescent Med J* 16:e11794. doi: 10.5812/ircmj.11794
- Nakhostin-Ansari, A., Sherafati, A., Aghajani, F., Khonji, M. S., Aghajani, R., and Shahmansouri, N. (2020). Depression and anxiety among Iranian medical students during COVID-19 pandemic. *Iran. J. Psychiatry* 15, 228–235. doi: 10.18502/ijps.v15i3.3815
- Oh, C.-M., Kim, H. Y., Na, H. K., Cho, K. H., and Chu, M. K. (2019). The effect of anxiety and depression on sleep quality of individuals with high risk for insomnia: a population-based study. *Front. Neurol.* 10:849. doi: 10.3389/fneur.2019.00849
- Örücü, M. Ç., and Demir, A. (2009). Psychometric evaluation of perceived stress scale for Turkish university students. *Stress Health* 25, 103–109. doi: 10.1080/09638237.2017.1417566
- Ozcan, H., Yucel, A., Avşar, U., Çankaya, E., Yucel, N., Gözübüyük, H., et al. (2015). Kidney transplantation is superior to hemodialysis and peritoneal dialysis in terms of cognitive function, anxiety, and depression symptoms in chronic kidney disease. *Transplant Proc* 47, 1348–1351. doi: 10.1016/j.transproceed.2015.04.032
- Pappa, S., Chen, J., Barnett, J., Chang, A., Dong, R. K., Xu, W., et al. (2022). A systematic review and meta-analysis of the mental health symptoms during the Covid-19 pandemic in Southeast Asia. *Psychiatry Clin. Neurosci.* 76, 41–50. doi: 10.1111/pcn.13306
- Politi, P., Piccinelli, M., and Wilkinson, G. (1994). Reliability, validity and factor structure of the 12-item general health questionnaire among young males in Italy. *Acta Psychiatr. Scand.* 90, 432–437. doi: 10.1111/j.1600-0447.1994.tb01620.x
- Proto, E., and Quintana-Domeque, C. (2021). COVID-19 and mental health deterioration by ethnicity and gender in the UK. *PLoS One* 16:e0244419. doi: 10.1371/journal.pone.0244419

- Revelle, W. R. (2017). *Psych: Procedures for Personality and Psychological Research [Computer software]*. Comprehensive R Archive Network.
- Roehrig, B., Hoeffken, K., Pientka, L., and Wedding, U. (2007). How many and which items of activities of daily living (ADL) and instrumental activities of daily living (IADL) are necessary for screening. *Crit. Rev. Oncol. Hematol.* 62, 164–171. doi: 10.1016/j.critrevonc.2006.10.001
- Rommel, M., Braehler, E., Roth, M., and Glaesmer, H. (2013). What is the general health Questionnaire-12 assessing?: dimensionality and psychometric properties of the general health Questionnaire-12 in a large scale German population sample. *Compr. Psychiatry* 54, 406–413. doi: 10.1016/j.comppsych.2012.10.010
- Santomauro, D. F., Herrera, A. M. M., Shadid, J., Zheng, P., Ashbaugh, C., Pigott, D. M., et al. (2021). Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *Lancet* 398, 1700–1712. doi: 10.1016/S0140-6736(21)02143-7
- Satorra, A., and Bentler, P. M. (2010). Ensuring Positiveness of the scaled difference chi-square test statistic. *Psychometrika* 75, 243–248. doi: 10.1007/s11336-009-9135-y
- Schnitz, N., Kruse, J., and Tress, W. (1999). Psychometric properties of the general health questionnaire (GHQ-12) in a German primary care sample. *Acta Psychiatr. Scand.* 100, 462–468. doi: 10.1111/j.1600-0447.1999.tb10898.x
- Shahriarirad, R., Erfani, A., Ranjbar, K., Bazrafshan, A., and Mirahmadzadeh, A. (2021). The mental health impact of COVID-19 outbreak: a Nationwide survey in Iran. *Int. J. Ment. Heal. Syst.* 15:19. doi: 10.1186/s13033-021-00445-3
- Sharifi, F., Alizadeh-Khoei, M., Saghebi, H., Angooti-Oshnari, L., Fadaee, S., Hormozi, S., et al. (2018). Validation study of ADL-Katz scale in the Iranian elderly nursing homes. *Ageing Int.* 43, 508–523. doi: 10.1007/s12126-017-9314-6
- Shojaei, S. F., and Masoumi, R. (2020). The importance of mental health training for psychologists in COVID-19 outbreak. *Middle East J Rehabil Health Stud* 7:e102846. doi: 10.5812/mejrh.102846
- Stuart, A., Katz, D., Stevenson, C., Gooch, D., Harkin, L., Bennasar, M., et al. (2022). Loneliness in older people and COVID-19: applying the social identity approach to digital intervention design. *Comput Hum Behav Rep* 6:100179. doi: 10.31234/osf.io/qk9hb
- Tang, N. K., Fiecas, M., Afolalu, E. F., and Wolke, D. (2017). Changes in sleep duration, quality, and medication use are prospectively associated with health and well-being: analysis of the UK household longitudinal study. *Sleep* 40. doi: 10.1093/sleep/zsw079
- Team, R. C. (2013). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. Available at: <http://www.R-project.org/> (Accessed April 23, 2022).
- Thielmann, B., Schierholz, R. S., and Böckelmann, I. (2021). Subjective and objective consequences of stress in subjects with subjectively different sleep quality—a cross-sectional study. *Int. J. Environ. Res. Public Health* 18:9990. doi: 10.3390/ijerph18199990
- Torales, J., Ríos-González, C., Barrios, I., O'Higgins, M., González, I., García, O., et al. (2020). Self-perceived stress during the quarantine of COVID-19 pandemic in Paraguay: an exploratory survey. *Front. Psych.* 11:558691. doi: 10.3389/fpsyg.2020.558691
- Varman, P. A., Ananthi, P., and Sugirda, P. (2022). Evaluation of perceived stress and coronavirus-specific anxiety among hospitalized COVID-19 patients—a cross-sectional study. *Asian J Med Sci* 13, 18–22. doi: 10.3126/ajms.v13i4.42132
- Wang, S., Li, B., Ungvari, G. S., Ng, C. H., Chiu, H. F., Kou, C., et al. (2016). Poor mental health status and its associations with demographic characteristics and chronic diseases in Chinese elderly. *Soc. Psychiatry Psychiatr. Epidemiol.* 51, 1449–1455. doi: 10.1007/s00127-016-1271-y
- WHO (2020). *Mental Health and Psychosocial Considerations during the COVID-19 Outbreak, 18 March 2020*. World Health Organization. Available at: <https://apps.who.int/iris/handle/10665/331490>
- Xiong, W., Liu, H., Gong, P., Wang, Q., Ren, Z., He, M., et al. (2019). Relationships of coping styles and sleep quality with anxiety symptoms among Chinese adolescents: a cross-sectional study. *J. Affect. Disord.* 257, 108–115. doi: 10.1016/j.jad.2019.07.032
- Zandifar, A., and Badrfam, R. (2020). Fighting COVID-19 in Iran; economic challenges ahead. *Arch. Iran. Med.* 23:284. doi: 10.34172/aim.2020.14
- Zhang, S. X., Miller, S. O., Xu, W., Yin, A., Chen, B. Z., Delios, A., et al. (2022). Meta-analytic evidence of depression and anxiety in Eastern Europe during the COVID-19 pandemic. *Eur. J. Psychotraumatol.* 13:2000132. doi: 10.1080/20008198.2021.2000132
- Zhong, X., Jin, X., Yan, L., Yang, L., Long, H., Wang, J., et al. (2021). Reliability and validity of General Health Questionnaire-12 in Chinese dental healthcare workers during the COVID-19 pandemic. *Front. Psych.* 12:792838. doi: 10.3389/fpsyg.2021.792838
- Zumbo, B. D., Gadermann, A. M., and Zeisser, C. J. O. M. A. S. M. (2007). Ordinal versions of coefficients alpha and theta for Likert rating scales. *J. Mod. Appl. Stat. Methods* 6, 21–29. doi: 10.22237/jmasm/1177992180